

LRPPRC Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21869c

Product Information

Application	WB, E
Primary Accession	P42704
Other Accession	Q6PB66 , Q5SGE0
Reactivity	Human, Rat, Mouse
Predicted	Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53981
Calculated MW	157905

Additional Information

Gene ID	10128
Other Names	Leucine-rich PPR motif-containing protein, mitochondrial, 130 kDa leucine-rich protein, LRP 130, GP130, LRPPRC, LRP130
Target/Specificity	This LRPPRC antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 847-876 amino acids from the Central region of human LRPPRC.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	LRPPRC Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	LRPPRC
Synonyms	LRP130
Function	May play a role in RNA metabolism in both nuclei and mitochondria. In the

nucleus binds to HNRPA1-associated poly(A) mRNAs and is part of nmRNP complexes at late stages of mRNA maturation which are possibly associated with nuclear mRNA export. Positively modulates nuclear export of mRNAs containing the EIF4E sensitivity element (4ESE) by binding simultaneously to both EIF4E and the 4ESE and acting as a platform for assembly for the RNA export complex (PubMed:[19262567](#), PubMed:[28325843](#)). Also binds to exportin XPO1/CRM1 to engage the nuclear pore and traffic the bound mRNAs to the cytoplasm (PubMed:[28325843](#)). May bind mature mRNA in the nucleus outer membrane. In mitochondria binds to poly(A) mRNA. Plays a role in translation or stability of mitochondrially encoded cytochrome c oxidase (COX) subunits. May be involved in transcription regulation. Cooperates with PPARGC1A to regulate certain mitochondrially encoded genes and gluconeogenic genes and may regulate docking of PPARGC1A to transcription factors. Seems to be involved in the transcription regulation of the multidrug-related genes MDR1 and MVP. Part of a nuclear factor that binds to the invMED1 element of MDR1 and MVP gene promoters. Binds single-stranded DNA (By similarity). Required for maintaining mitochondrial potential (PubMed:[23822101](#)). Suppresses the initiation of basal levels of autophagy and mitophagy by sustaining BCL2 levels (PubMed:[23822101](#)).

Cellular Location

Mitochondrion. Nucleus Nucleus, nucleoplasm. Nucleus inner membrane. Nucleus outer membrane Note=Seems to be predominantly mitochondrial

Tissue Location

Expressed ubiquitously. Expression is highest in heart, skeletal muscle, kidney and liver, intermediate in brain, non- mucosal colon, spleen and placenta, and lowest in small intestine, thymus, lung and peripheral blood leukocytes

Background


May play a role in RNA metabolism in both nuclei and mitochondria. In the nucleus binds to HNRPA1-associated poly(A) mRNAs and is part of nmRNP complexes at late stages of mRNA maturation which are possibly associated with nuclear mRNA export. May bind mature mRNA in the nucleus outer membrane. In mitochondria binds to poly(A) mRNA. Plays a role in translation or stability of mitochondrially encoded cytochrome c oxidase (COX) subunits. May be involved in transcription regulation. Cooperates with PPARGC1A to regulate certain mitochondrially encoded genes and gluconeogenic genes and may regulate docking of PPARGC1A to transcription factors. Seems to be involved in the transcription regulation of the multidrug-related genes MDR1 and MVP. Part of a nuclear factor that binds to the invMED1 element of MDR1 and MVP gene promoters. Binds single-stranded DNA (By similarity).

References

Xu F.,et al.Biochem. J. 382:331-336(2004).
 Ota T.,et al.Nat. Genet. 36:40-45(2004).
 Hillier L.W.,et al.Nature 434:724-731(2005).
 Hou J.,et al.In Vitro Cell. Dev. Biol. Anim. 30A:111-114(1994).
 Bienvenut W.V.,et al.Submitted (JUL-2007) to UniProtKB.

Images

All lanes : Anti-LRPPRC Antibody (Center) at 1:2000
 dilution Lane 1: 293T/17 whole cell lysate Lane 2: HeLa
 whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4:
 mouse liver lysate Lysates/proteins at 20 µg per lane.
 Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase
 conjugated at 1/10000 dilution. Predicted band size : 158



kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.