

NAT8L Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21925a

Product Information

Application	WB, IHC-P-Leica, E
Primary Accession	Q8N9F0
Other Accession	Q3UGX3 , D3ZVU9
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB55378
Calculated MW	32837

Additional Information

Gene ID	339983
Other Names	N-acetylaspartate synthetase, NAA synthetase, 2.3.1.17, Camello-like protein 3, N-acetyltransferase 8-like protein, NAT8L, CML3
Target/Specificity	This NAT8L antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 59-93 amino acids from human NAT8L.
Dilution	WB~~1:2000 IHC-P-Leica~~1:250 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NAT8L Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NAT8L (HGNC:26742)
Synonyms	CML3
Function	Catalyzes the synthesis of N-acetylaspartate acid (NAA) from L-aspartate

and acetyl-CoA (PubMed:[19524112](#), PubMed:[19807691](#), PubMed:[20385109](#)). Promotes dopamine uptake by regulating TNF expression (By similarity). Attenuates methamphetamine-induced inhibition of dopamine uptake (PubMed:[20385109](#)).

Cellular Location

Cytoplasm. Microsome membrane {ECO:0000250|UniProtKB:D3ZVU9}; Single-pass membrane protein. Mitochondrion membrane; Single-pass membrane protein. Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q3UGX3}; Single-pass membrane protein. Note=Its enzymatic activity contribution is quantitatively larger in mitochondrial compartment than in extramitochondrial compartment.

Tissue Location

Expressed in brain..


Background

Plays a role in the regulation of lipogenesis by producing N-acetylaspartate acid (NAA), a brain-specific metabolite. NAA occurs in high concentration in brain and its hydrolysis plays a significant part in the maintenance of intact white matter. Promotes dopamine uptake by regulating TNF-alpha expression. Attenuates methamphetamine-induced inhibition of dopamine uptake.


References

- Hillier L.W.,et al.Nature 434:724-731(2005).
Brandenberger R.,et al.Nat. Biotechnol. 22:707-716(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Popsueva A.E.,et al.Dev. Biol. 234:483-496(2001).
Arun P.,et al.Neurochem. Int. 55:219-225(2009).

Images



Immunohistochemical analysis of AP21925A on paraffin-embedded human kidney tissue was performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:250) for 15min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.



Anti-NAT8L Antibody (N-Term) at 1:1000 dilution + U-87 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.