

PC2 (CBX4) Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2514a

Product Information

Application	WB, IHC-P, E
Primary Accession	O00257
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	61368
Antigen Region	80-110

Additional Information

Gene ID	8535
Other Names	E3 SUMO-protein ligase CBX4, 632-, Chromobox protein homolog 4, Polycomb 2 homolog, Pc2, hPc2, CBX4
Target/Specificity	This PC2 (CBX4) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 80-110 amino acids from the N-terminal region of human PC2 (CBX4).
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PC2 (CBX4) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CBX4
Function	E3 SUMO-protein ligase that catalyzes sumoylation of target proteins by promoting the transfer of SUMO from the E2 enzyme to the substrate (PubMed: 12679040 , PubMed: 22467880 , PubMed: 22825850). Also acts as a histone reader, which specifically recognizes and binds histone H3 trimethylated at 'Lys-9' and 'Lys-27' (H3K9me3 and H3K27me3, respectively)

via its chromo domain (By similarity). Catalyzes sumoylation of HNRNPK, a p53/TP53 transcriptional coactivator, hence indirectly regulates p53/TP53 transcriptional activation resulting in p21/CDKN1A expression (PubMed:[22825850](#)). Acts as a regulator of brown adipocyte differentiation by mediating sumoylation of PRDM16, thereby preventing PRDM16 ubiquitination and degradation (By similarity). Monosumoylates ZNF131 (PubMed:[22467880](#)). Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development (PubMed:[12167701](#), PubMed:[19636380](#), PubMed:[21282530](#)). PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:[12167701](#), PubMed:[19636380](#), PubMed:[21282530](#)). Plays a role in the lineage differentiation of the germ layers in embryonic development (By similarity).

Cellular Location

Nucleus. Nucleus speckle. Note=Localization to nuclear polycomb bodies is required for ZNF131 sumoylation (PubMed:[22467880](#)). Localized in distinct foci on chromatin (PubMed:[18927235](#))

Tissue Location

Ubiquitous.


Background

CBX4 is a member of Drosophila Polycomb group gene family. The polycomb group (PcG) genes are essential for maintenance of proper expression patterns of developmental master regulators; changes in expression of PcG proteins have been associated with cancer. CBX4 is a part of the cellular memory system responsible for the inheritance of gene activity by progeny cells. It participates in maintaining the transcriptionally repressive state of genes. CBX4 is part of a complex that acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. CBX4 is an E3 SUMO-protein ligase which facilitates SUMO1 conjugation by UBE2I.

References


Kagey, M.H., et al., Cell 113(1):127-137 (2003).
Satijn, D.P., et al., Mol. Cell. Biol. 17(10):6076-6086 (1997).

Images



The anti-CBX4 N-term Pab (Cat. #AP2514a) is used in Western blot to detect CBX4 in mouse kidney tissue lysate.

Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use



of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Citations

- [HSP70-Hrd1 axis precludes the oncorepressor potential of N-terminal misfolded Blimp-1s in lymphoma cells.](#)
- [Human Polycomb protein 2 promotes \$\alpha\$ -synuclein aggregate formation through covalent SUMOylation.](#)
- [The SUMO E3 ligase activity of Pc2 is coordinated through a SUMO interaction motif.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.