

# Phospho-H3(S10) Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP3003a

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">P68431</a>
<b>Other Accession</b>	<a href="#">P61830</a> , <a href="#">P02299</a> , <a href="#">P08898</a> , <a href="#">P02302</a> , <a href="#">P02301</a> , <a href="#">Q6NXT2</a> , <a href="#">A5PK61</a> , <a href="#">Q6PI79</a> , <a href="#">P84245</a> , <a href="#">P84246</a> , <a href="#">Q71LE2</a> , <a href="#">P84244</a> , <a href="#">P84243</a> , <a href="#">P84249</a> , <a href="#">Q6PI20</a> , <a href="#">P84247</a> , <a href="#">Q5E9F8</a> , <a href="#">Q27532</a> , <a href="#">Q9U281</a> , <a href="#">Q10453</a> , <a href="#">P84233</a> , <a href="#">P84228</a> , <a href="#">Q71DI3</a> , <a href="#">Q4QRF4</a> , <a href="#">P84229</a> , <a href="#">P84227</a> , <a href="#">Q6LED0</a> , <a href="#">P68433</a> , <a href="#">P68432</a> , <a href="#">Q16695</a> , <a href="#">Q71DJ3</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Bovine, Mouse, Rat, Chicken, Zebrafish, Xenopus, C.Elegans, Drosophila, Pig, Rabbit, Yeast
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB5807
<b>Calculated MW</b>	15404

## Additional Information

---

<b>Gene ID</b>	8350;8351;8352;8353;8354;8355;8356;8357;8358;8968
<b>Other Names</b>	Histone H31, Histone H3/a, Histone H3/b, Histone H3/c, Histone H3/d, Histone H3/f, Histone H3/h, Histone H3/i, Histone H3/j, Histone H3/k, Histone H3/l, HIST1H3A, H3FA
<b>Target/Specificity</b>	This H3 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S10 of human H3.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Phospho-H3(S10) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	H3C1 ( <a href="#">HGNC:4766</a> )
<b>Synonyms</b>	H3FA, HIST1H3A
<b>Function</b>	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
<b>Cellular Location</b>	Nucleus. Chromosome.


## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. The gene for this protein is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. The gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

## References


- Lusic, M., et al., *EMBO J.* 22(24):6550-6561 (2003).  
Deng, L., et al., *Virology* 289(2):312-326 (2001).  
Deng, L., et al., *Virology* 277(2):278-295 (2000).  
El Kharroubi, A., et al., *Mol. Cell. Biol.* 18(5):2535-2544 (1998).  
Albig, W., et al., *Hum. Genet.* 101(3):284-294 (1997).

## Images



Western blot analysis of anti-Phospho-H3-pS10 Pab (Cat. #AP3003a) in CEM cell line lysate (35ug/lane). Phospho-H3-pS10(arrow) was detected using the purified Pab.

Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.