

Phospho-Dnmt1(S154) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3534a

Product Information

Application	DB, E
Primary Accession	P26358
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB18429
Calculated MW	183165

Additional Information

Gene ID	1786
Other Names	DNA (cytosine-5)-methyltransferase 1, Dnmt1, CXXC-type zinc finger protein 9, DNA methyltransferase HsaI, DNA MTase HsaI, MHsaI, MCMT, DNMT1, AIM, CXXC9, DNMT
Target/Specificity	This Dnmt1 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S154 of human Dnmt1.
Dilution	DB~1:500 E~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-Dnmt1(S154) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DNMT1 {ECO:0000303 Ref.3, ECO:0000312 HGNC:HGNC:2976}
Function	DNA methyltransferase that methylates CpG residues (PubMed: 17200670 , PubMed: 18754681 , PubMed: 21745816 , PubMed: 26070743). Preferentially methylates hemimethylated DNA (PubMed: 21745816 , PubMed: 26070743). Associates with DNA replication sites in S phase maintaining the methylation

pattern in the newly synthesized strand, that is essential for epigenetic inheritance (PubMed:[17200670](#), PubMed:[21745816](#)). Associates with chromatin during G2 and M phases to maintain DNA methylation independently of replication (PubMed:[21745816](#)). It is responsible for maintaining methylation patterns established in development (PubMed:[21745816](#)). DNA methylation is coordinated with methylation of histones (PubMed:[16357870](#)). Mediates transcriptional repression by direct binding to HDAC2 (PubMed:[10888872](#)). In association with DNMT3B and via the recruitment of CTCFL/BORIS, involved in activation of BAG1 gene expression by modulating dimethylation of promoter histone H3 at H3K4 and H3K9 (PubMed:[18413740](#)). Probably forms a corepressor complex required for activated KRAS-mediated promoter hypermethylation and transcriptional silencing of tumor suppressor genes (TSGs) or other tumor-related genes in colorectal cancer (CRC) cells (PubMed:[24623306](#)). Also required to maintain a transcriptionally repressive state of genes in undifferentiated embryonic stem cells (ESCs) (PubMed:[24623306](#)). Associates at promoter regions of tumor suppressor genes (TSGs) leading to their gene silencing (PubMed:[24623306](#)).

Cellular Location

Nucleus. Chromosome Note=Associates with replication foci during S-phase: recruited to hemimethylated DNA sites via its RFTS domain, which specifically recognizes and binds histone H3 ubiquitinated at 'Lys-14', 'Lys-18' and 'Lys-23' (H3K14ub, H3K18ub and H3K23ub, respectively) (PubMed:[29053958](#)). Localized to the perinucleolar region (PubMed:[24492612](#)).

Tissue Location

Ubiquitous; highly expressed in fetal tissues, heart, kidney, placenta, peripheral blood mononuclear cells, and expressed at lower levels in spleen, lung, brain, small intestine, colon, liver, and skeletal muscle. Isoform 2 is less expressed than isoform 1.


Background

DNA (cytosine-5-)-methyltransferase 1 has a role in the establishment and regulation of tissue-specific patterns of methylated cytosine residues. Aberrant methylation patterns are associated with certain human tumors and developmental abnormalities.

References

- Liao,X., Int. J. Cancer 123 (2), 296-302 (2008)
Leng,S., (er) Carcinogenesis (2008) In press
Dion,V., Hum. Mol. Genet. 17 (9), 1306-1317 (2008)

Images



Dot blot analysis of anti-Phospho-Dnmt1-pS154 Antibody (Cat.#AP3534a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.