

# Cullin 2 Antibody

Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP50189

## Product Information

---

Application	WB, IF, IHC
Primary Accession	<a href="#">Q13617</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Calculated MW	86983

## Additional Information

---

Gene ID	8453
Other Names	Cullin-2, CUL-2, CUL2
Dilution	WB~~ 1:1000 IF~~1:100 IHC~~1:50-1:100
Format	Rabbit IgG in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

## Protein Information

---

Name	CUL2 ( <a href="#">HGNC:2552</a> )
Function	<p>Core component of multiple cullin-RING-based ECS (ElonginB/C-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of target proteins (PubMed:<a href="#">11384984</a>, PubMed:<a href="#">26138980</a>, PubMed:<a href="#">29775578</a>, PubMed:<a href="#">29779948</a>, PubMed:<a href="#">37844242</a>, PubMed:<a href="#">38326650</a>). CUL2 serves as a rigid scaffold in the complex and may contribute to catalysis through positioning of the substrate and the E2 ubiquitin-conjugating enzyme (PubMed:<a href="#">10973499</a>, PubMed:<a href="#">11384984</a>, PubMed:<a href="#">12609982</a>, PubMed:<a href="#">24076655</a>, PubMed:<a href="#">9122164</a>, PubMed:<a href="#">37844242</a>, PubMed:<a href="#">38326650</a>). The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (PubMed:<a href="#">12609982</a>, PubMed:<a href="#">24076655</a>, PubMed:<a href="#">27565346</a>, PubMed:<a href="#">38326650</a>). The functional specificity of the ECS complex depends on the substrate recognition component (PubMed:<a href="#">10973499</a>, PubMed:<a href="#">26138980</a>, PubMed:<a href="#">29775578</a>, PubMed:<a href="#">29779948</a>, PubMed:<a href="#">9122164</a>, PubMed:<a href="#">38326650</a>). ECS(VHL) mediates the ubiquitination of hypoxia-inducible factor (HIF) (PubMed:<a href="#">10973499</a>, PubMed:<a href="#">9122164</a>). A number of ECS complexes (containing either KLHDC2, KLHDC3, KLHDC10, APPBP2,</p>

FEM1A, FEM1B or FEM1C as substrate-recognition component) are part of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiquitination and degradation (PubMed:[26138980](#), PubMed:[29775578](#), PubMed:[29779948](#), PubMed:[37844242](#)). ECS complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins (PubMed:[27565346](#)). ECS(LRR1) ubiquitinates MCM7 and promotes CMG replisome disassembly by VCP and chromatin extraction during S- phase (By similarity).

**Cellular Location** Nucleus {ECO:0000250|UniProtKB:Q9D4H8}.


## Background

Core component of multiple cullin-RING-based ECS (ElonginB/C-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of target proteins. May serve as a rigid scaffold in the complex and may contribute to catalysis through positioning of the substrate and the ubiquitin- conjugating enzyme. The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (By similarity). The functional specificity of the ECS complex depends on the substrate recognition component. ECS(VHL) mediates the ubiquitination of hypoxia-inducible factor (HIF).


## References

Pause A.,et al.Proc. Natl. Acad. Sci. U.S.A. 94:2156-2161(1997).  
Wada H.,et al.Biochem. Biophys. Res. Commun. 257:100-105(1999).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Deloukas P.,et al.Nature 429:375-381(2004).  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

## Images





Western blot analysis of lysate from MDA-MB-231 cell line, using Cullin 2 Antibody (C0163). C0163 was diluted at 1:1000. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



Immunofluorescence analysis of NIH/3T3 cells, using Cullin 2 antibody .

Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using Cullin 2 antibody .



Western blot analysis of extracts from HepG2 cells (Lane 2), using Cullin 2 Antibody. The lane on the left is treated with synthesized peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.