

# Cytochrome P450 39A1 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP50820

#### **Product Information**

Application WB
Primary Accession Q9NYL5
Reactivity Human
Host Rabbit
Clonality polyclonal
Calculated MW 54116

## **Additional Information**

**Gene ID** 51302

Other Names 24-hydroxycholesterol 7-alpha-hydroxylase, Cytochrome P450 39A1,

hCYP39A1, Oxysterol 7-alpha-hydroxylase, CYP39A1

**Dilution** WB~~ 1:1000

Format Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4,

150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions -20°C

#### **Protein Information**

Name CYP39A1 {ECO:0000303 | PubMed:25201972,

ECO:0000312 | HGNC:HGNC:17449}

**Function** A cytochrome P450 monooxygenase involved in neural cholesterol clearance

through bile acid synthesis (PubMed:10748047, PubMed:25201972). Catalyzes 7-alpha hydroxylation of (24S)- hydroxycholesterol, a neural oxysterol that is metabolized to bile acids in the liver (PubMed:10748047, PubMed:25201972). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons

provided by NADPH via cytochrome P450 reductase (CPR;

NADPH-ferrihemoprotein reductase) (PubMed: 10748047, PubMed: 25201972).

**Cellular Location** Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q64654};

Multi-pass membrane protein. Microsome membrane

{ECO:0000250|UniProtKB:Q64654}; Multi-pass membrane protein

Tissue Location Liver specific...


## **Background**

Involved in the bile acid metabolism. Has a preference for 24-hydroxycholesterol, and converts it into a 7-alpha- hydroxylated product.

## References

Li-Hawkins J., et al.J. Biol. Chem. 275:16543-16549(2000). Ota T., et al. Nat. Genet. 36:40-45(2004). Mungall A.J., et al. Nature 425:805-811(2003).

# **Images**



Western blot analysis of lysate from human liver tissue lysate, using Cytochrome P450 39A1 Antibody, was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.