

# **IKZF2** Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP50853

### **Product Information**

Application WB
Primary Accession Q9UKS7

**Reactivity** Human, Mouse, Rat

HostRabbitClonalitypolyclonalCalculated MW59574

### **Additional Information**

**Gene ID** 22807

Other Names Zinc finger protein Helios, Ikaros family zinc finger protein 2, IKZF2, HELIOS,

ZNFN1A2

**Dilution** WB~~ 1:1000

Format Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4,

150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions -20°C

## **Protein Information**

Name IKZF2

**Synonyms** HELIOS, ZNFN1A2

**Function** Transcriptional regulator required for outer hair cells (OHC) maturation and,

consequently, for hearing.

**Cellular Location** Nucleus {ECO:0000250 | UniProtKB:P81183}. Note=Associates with Ikaros at

centromeric heterochromatin in T-cell nuclei.

{ECO:0000250 | UniProtKB:P81183}


# **Background**

Associates with Ikaros at centromeric heterochromatin.

## References

Hosokawa Y., et al. Immunogenetics 50:106-108(1999). Perdomo J., et al. J. Biol. Chem. 275:38347-38354(2000). Mayya V., et al. Sci. Signal. 2:RA46-RA46(2009). Choudhary C., et al. Science 325:834-840(2009).

# **Images**



Western blot analysis of lysate from KG-1 cell line, using IKZF2 Antibody, was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35 ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.