

Aquaporin 3 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50990

Product Information

Application	WB, IHC-P
Primary Accession	Q92482
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31544

Additional Information

Gene ID	360
Other Names	Aquaporin-3, AQP-3, Aquaglyceroporin-3, AQP3
Dilution	WB~~1:1000 IHC-P~~N/A
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	AQP3 {ECO:0000303 PubMed:7558005, ECO:0000312 HGNC:HGNC:636}
Function	Aquaglyceroporins form homotetrameric transmembrane channels, with each monomer independently mediating glycerol and water transport across the plasma membrane along their osmotic gradient (PubMed: 12239222 , PubMed: 30420639). Could also be permeable to urea (By similarity). Also participates in cell permeability to H2O2 and H2O2- mediated signaling (PubMed: 20724658). In skin, transports glycerol to the epidermis and stratum corneum, where it maintains hydration, elasticity, and supports lipid biosynthesis for barrier repair (By similarity). In kidney, contributes to the reabsorption of water, helping the body maintain proper fluid balance (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:O14520}. Basolateral cell membrane {ECO:0000250 UniProtKB:P47862}; Multi-pass membrane protein {ECO:0000250 UniProtKB:O14520}
Tissue Location	Widely expressed in epithelial cells of kidney (collecting ducts) and airways, in keratinocytes, immature dendritic cells and erythrocytes. Isoform 2 is not detectable in erythrocytes at the protein level

Background

Water channel required to promote glycerol permeability and water transport across cell membranes. Acts as a glycerol transporter in skin and plays an important role in regulating SC (stratum corneum) and epidermal glycerol content. Involved in skin hydration, wound healing, and tumorigenesis. Provides kidney medullary collecting duct with high permeability to water, thereby permitting water to move in the direction of an osmotic gradient. Slightly permeable to urea and may function as a water and urea exit mechanism in antidiuresis in collecting duct cells. It may play an important role in gastrointestinal tract water transport and in glycerol metabolism (By similarity).

References

- Ishibashi K.,et al.Genomics 27:352-354(1995).
Ishibashi K.,et al.Submitted (OCT-1996) to the EMBL/GenBank/DDBJ databases.
Roudier N.,et al.J. Biol. Chem. 277:45854-45859(2002).
Halleck A.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.