

Cytochrome P450 2J2 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51147

Product Information

Application	WB
Primary Accession	P51589
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57611

Additional Information

Gene ID	1573
Other Names	Cytochrome P450 2J2, Arachidonic acid epoxygenase, CYP11J2, CYP2J2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Cytochrome P450 2J2. The exact sequence is proprietary.
Dilution	WB~~ 1:2000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CYP2J2 {ECO:0000303 PubMed:19737933, ECO:0000312 HGNC:HGNC:2634}
Function	<p>A cytochrome P450 monooxygenase involved in the metabolism of polyunsaturated fatty acids (PUFA) in the cardiovascular system (PubMed:19965576, PubMed:8631948). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:19965576, PubMed:8631948). Catalyzes the epoxidation of double bonds of PUFA (PubMed:19965576, PubMed:8631948). Converts arachidonic acid to four regioisomeric epoxygenic acids (EpETs), likely playing a major role in the epoxidation of endogenous cardiac arachidonic acid pools (PubMed:8631948). In endothelial cells, participates in eicosanoids metabolism by converting hydroperoxide species into hydroxy epoxy metabolites. In combination with 15- lipoxygenase metabolizes arachidonic acid and converts hydroperoxyicosatetraenoates (HpETs) into hydroxy epoxy eicosatrienoates (HEETs), which are precursors of vasodilatory trihydroxyicosatrienoic acids (THETAs). This hydroperoxide isomerase activity</p>

is NADPH- and O₂-independent (PubMed:[19737933](#)). Catalyzes the monooxygenation of a various xenobiotics, such as danazol, amiodarone, terfenadine, astemizole, thioridazine, tamoxifen, cyclosporin A and nabumetone (PubMed:[19923256](#)). Catalyzes hydroxylation of the anthelmintics albendazole and fenbendazole (PubMed:[23959307](#)). Catalyzes the sulfoxidation of fenbedazole (PubMed:[19923256](#)).

Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

Tissue Location

Highly expressed in heart, present at lower levels in liver, kidney and skeletal muscle (at protein level)

Background

This enzyme metabolizes arachidonic acid predominantly via a NADPH-dependent olefin epoxidation to all four regioisomeric cis-epoxyeicosatrienoic acids. One of the predominant enzymes responsible for the epoxidation of endogenous cardiac arachidonic acid pools.


References

Wu S.,et al.J. Biol. Chem. 271:3460-3468(1996).

Wu S.,et al.Submitted (JAN-2002) to the EMBL/GenBank/DDBJ databases.

King L.M.,et al.Mol. Pharmacol. 61:840-852(2002).

Images



All lanes : Anti-Cytochrome P450 2J2 Antibody at 1:2000 dilution Lane 1: A549 whole cell lysates Lane 2: A431 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 58 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.