

# DNAJB11 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51166

## Product Information

---

Application	WB
Primary Accession	<a href="#">Q9UBS4</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40514

## Additional Information

---

Gene ID	51726
Other Names	DnaJ homolog subfamily B member 11, APOBEC1-binding protein 2, ABBP-2, DnaJ protein homolog 9, ER-associated DNAJ, ER-associated Hsp40 co-chaperone, Endoplasmic reticulum DNA J domain-containing protein 3, ER-resident protein ERdj3, ERdj3, ERj3p, HEDJ, Human DnaJ protein 9, hDj-9, PWP1-interacting protein 4, DNAJB11, EDJ, ERJ3, HDJ9
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human DNAJB11. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	DNAJB11
Synonyms	EDJ, ERJ3, HDJ9
Function	As a co-chaperone for HSPA5 it is required for proper folding, trafficking or degradation of proteins (PubMed: <a href="#">10827079</a> , PubMed: <a href="#">15525676</a> , PubMed: <a href="#">29706351</a> ). Binds directly to both unfolded proteins that are substrates for ERAD and nascent unfolded peptide chains, but dissociates from the HSPA5-unfolded protein complex before folding is completed (PubMed: <a href="#">15525676</a> ). May help recruiting HSPA5 and other chaperones to the substrate. Stimulates HSPA5 ATPase activity (PubMed: <a href="#">10827079</a> ). It is necessary for maturation and correct trafficking of PKD1 (PubMed: <a href="#">29706351</a> ).
Cellular Location	Endoplasmic reticulum lumen Note=Associated with the ER membrane in a

C-terminally epitope-tagged construct

## Tissue Location

Widely expressed.

## Background

Serves as a co-chaperone for HSPA5. Binds directly to both unfolded proteins that are substrates for ERAD and nascent unfolded peptide chains, but dissociates from the HSPA5-unfolded protein complex before folding is completed. May help recruiting HSPA5 and other chaperones to the substrate. Stimulates HSPA5 ATPase activity.

## References

Ohtsuka K.,et al.Cell Stress Chaperones 5:98-112(2000).


Yu M.,et al.J. Biol. Chem. 275:24984-24992(2000).

Bies C.,et al.Biol. Chem. 385:389-395(2004).

Honore B.,et al.Submitted (JUN-2000) to the EMBL/GenBank/DDBJ databases.

Clark H.F.,et al.Genome Res. 13:2265-2270(2003).

## Images



All lanes : Anti-DNAJB11 Antibody at 1:1000 dilution Lane 1: HeLa whole cell lysates Lane 2: K562 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 41 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.