

MDA5 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51277

Product Information

Application WB, ICC, IHC-P

Primary Accession

Reactivity
Host
Clonality
Calculated MW

Q9BYX4
Human, Rat
Rabbit
Polyclonal
116689

Additional Information

Gene ID 64135

Other Names Interferon-induced helicase C domain-containing protein 1, Clinically

amyopathic dermatomyositis autoantigen 140 kDa, CADM-140 autoantigen, Helicase with 2 CARD domains, Helicard, Interferon-induced with helicase C domain protein 1, Melanoma differentiation-associated protein 5, MDA-5, Murabutide down-regulated protein, RIG-I-like receptor 2, RLR-2, RNA

helicase-DEAD box protein 116, IFIH1, MDA5, RH116

Target/Specificity KLH-conjugated synthetic peptide encompassing a sequence within the

C-term region of human MDA5. The exact sequence is proprietary.

Dilution WB~~1:1000 ICC~~N/A IHC-P~~N/A

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name IFIH1 (HGNC:18873)

Function Innate immune receptor which acts as a cytoplasmic sensor of viral nucleic

acids and plays a major role in sensing viral infection and in the activation of a cascade of antiviral responses including the induction of type I interferons and pro-inflammatory cytokines (PubMed:28594402, PubMed:32169843, PubMed:33727702). Its ligands include mRNA lacking 2'-O-methylation at their 5' cap and long-dsRNA (>1 kb in length) (PubMed:22160685). Upon ligand binding it associates with mitochondria antiviral signaling protein (MAVS/IPS1) which activates the IKK-related kinases: TBK1 and IKBKE which phosphorylate interferon regulatory factors: IRF3 and IRF7 which in turn activate transcription of antiviral immunological genes, including interferons (IFNs); IFN-alpha and IFN-beta. Responsible for detecting the Picornaviridae

family members such as encephalomyocarditis virus (EMCV), mengo encephalomyocarditis virus (ENMG), and rhinovirus (PubMed: 28606988). Detects coronavirus SARS-CoV-2 (PubMed: 33440148, PubMed: 33514628). Can also detect other viruses such as dengue virus (DENV), west Nile virus (WNV), and reovirus. Also involved in antiviral signaling in response to viruses containing a dsDNA genome, such as vaccinia virus. Plays an important role in amplifying innate immune signaling through recognition of RNA metabolites that are produced during virus infection by ribonuclease L (RNase L). May play an important role in enhancing natural killer cell function and may be involved in growth inhibition and apoptosis in several tumor cell lines.

Cellular Location Cytoplasm. Nucleus. Mitochondrion. Note=Upon viral RNA stimulation and

ISGylation, translocates from cytosol to mitochondrion. May be found in the

nucleus, during apoptosis

Tissue Location Widely expressed, at a low level. Expression is detected at slightly highest

levels in placenta, pancreas and spleen and at barely levels in detectable

brain, testis and lung

Background

Innate immune receptor which acts as a cytoplasmic sensor of viral nucleic acids and plays a major role in sensing viral infection and in the activation of a cascade of antiviral responses including the induction of type I interferons and proinflammatory cytokines. Its ligands include mRNA lacking 2'-O- methylation at their 5' cap and long-dsRNA (>1 kb in length). Upon ligand binding it associates with mitochondria antiviral signaling protein (MAVS/IPS1) which activates the IKK-related kinases: TBK1 and IKBKE which phosphorylate interferon regulatory factors: IRF3 and IRF7 which in turn activate transcription of antiviral immunological genes, including interferons (IFNs); IFN-alpha and IFN-beta. Responsible for detecting the Picornaviridae family members such as encephalomyocarditis virus (EMCV) and mengo encephalomyocarditis virus (ENMG). Can also detect other viruses such as dengue virus (DENV), west Nile virus (WNV), and reovirus. Also involved in antiviral signaling in response to viruses containing a dsDNA genome, such as vaccinia virus. Plays an important role in amplifying innate immune signaling through recognition of RNA metabolites that are produced during virus infection by ribonuclease L (RNase L). May play an important role in enhancing natural killer cell function and may be involved in growth inhibition and apoptosis in several tumor cell lines.

References

Kang D.-C., et al. Proc. Natl. Acad. Sci. U.S.A. 99:637-642(2002). Cocude C., et al. J. Gen. Virol. 84:3215-3225(2003). Ota T., et al. Nat. Genet. 36:40-45(2004). Kovacsovics M., et al. Curr. Biol. 12:838-843(2002). Kovacsovics M., et al. Curr. Biol. 12:1633-1633(2002).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.