

GANP Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51336

Product Information

Application	WB, IHC-P
Primary Accession	O60318
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	218405

Additional Information

Gene ID	8888
Other Names	Germinal-center associated nuclear protein, GANP, 80 kDa MCM3-associated protein, MCM3 acetylating protein, MCM3AP, 231-, MCM3 acetyltransferase, MCM3AP, GANP, KIAA0572, MAP80
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human GANP. The exact sequence is proprietary.
Dilution	WB~~1:1000 IHC-P~~N/A
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	MCM3AP
Synonyms	GANP, KIAA0572, MAP80
Function	[Isoform GANP]: As a component of the TREX-2 complex, involved in the export of mRNAs to the cytoplasm through the nuclear pores (PubMed: 20005110 , PubMed: 20384790 , PubMed: 22307388 , PubMed: 23591820). Through the acetylation of histones, affects the assembly of nucleosomes at immunoglobulin variable region genes and promotes the recruitment and positioning of transcription complex to favor DNA cytosine deaminase AICDA/AID targeting, hence promoting somatic hypermutations (PubMed: 23652018).
Cellular Location	[Isoform GANP]: Nucleus envelope. Nucleus, nuclear pore complex. Nucleus, nucleoplasm. Chromosome. Note=Predominantly located at the nuclear envelope, facing the nucleus interior (PubMed:20005110, PubMed:21195085,

PubMed:23591820). Localization at the nuclear pore complex requires NUP153, TPR and ALYREF/ALY (PubMed:22307388, PubMed:23591820). Also found associated with chromatin (PubMed:23652018). In B-cells, targeted to the immunoglobulin variable region genes (PubMed:23652018).

Tissue Location

Widely expressed (PubMed:11024281). Up-regulated in germinal center B-cells in tonsils (at protein level) (PubMed:11024281).

Background

Isoform GANP: Essential for the generation of high- affinity B-cells against T-cell-dependent antigens by affecting somatic hypermutation at the IgV-regions. May have stimulation- dependent DNA primase activity that would generate extra RNA primers in very rapidly proliferating cells and would support clonal expansion of differentiating B-cells (By similarity). Involved in the nuclear export of poly(A)-containing mRNAs by acting as a scaffold for the TREX-2 complex. The TREX-2 complex functions in docking export-competent ribonucleoprotein particles (mRNPs) to the nuclear entrance of the nuclear pore complex (nuclear basket). TREX-2 participates in mRNA export and accurate chromatin positioning in the nucleus by tethering genes to the nuclear periphery.

References

Abe E.,et al.Gene 255:219-227(2000).
Hattori M.,et al.Nature 405:311-319(2000).
Nagase T.,et al.DNA Res. 5:31-39(1998).
Takei Y.,et al.J. Biol. Chem. 273:22177-22180(1998).
Takei Y.,et al.EMBO Rep. 2:119-123(2001).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.