

# OCT1 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51439

## Product Information

---

Application	WB
Primary Accession	<a href="#">P14859</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	76472

## Additional Information

---

Gene ID	5451
Other Names	POU domain, class 2, transcription factor 1, NF-A1, Octamer-binding protein 1, Oct-1, Octamer-binding transcription factor 1, OTF-1, POU2F1, OCT1, OTF1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human 41913. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	POU2F1
Synonyms	OCT1, OTF1
Function	Transcription factor that binds to the octamer motif (5'- ATTTGCAT-3') and activates the promoters of the genes for some small nuclear RNAs (snRNA) and of genes such as those for histone H2B and immunoglobulins. Modulates transcription transactivation by NR3C1, AR and PGR.
Cellular Location	Nucleus.
Tissue Location	Ubiquitous. Isoform 2 is lymphocyte-specific.

## Background

---

Transcription factor that binds to the octamer motif (5'-ATTTGCAT-3') and activates the promoters of the genes for some small nuclear RNAs (snRNA) and of genes such as those for histone H2B and immunoglobulins. Modulates transcription transactivation by NR3C1, AR and PGR (By similarity). In case of human herpes simplex virus (HSV) infection, POU2F1 forms a multiprotein-DNA complex with the viral transactivator protein VP16 and HCFC1 thereby enabling the transcription of the viral immediate early genes.


## References

---

Sturm R.A.,et al.Genes Dev. 2:1582-1599(1988).  
Luchina N.N.,et al.Immunol. Lett. 85:237-241(2003).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Gregory S.G.,et al.Nature 441:315-321(2006).  
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

## Images

---



All lanes : Anti-OCT1 Antibody at 1:1000 dilution Lane 1: HeLa whole cell lysates Lane 2: K562 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 76 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.