

# AKR1A1 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51718

## Product Information

---

Application	WB
Primary Accession	<a href="#">P14550</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36573

## Additional Information

---

Gene ID	10327
Other Names	Alcohol dehydrogenase [NADP(+)], Aldehyde reductase, Aldo-keto reductase family 1 member A1, AKR1A1, ALDR1, ALR
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human AKR1A1. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	AKR1A1
Synonyms	ALDR1, ALR
Function	Catalyzes the NADPH-dependent reduction of a wide variety of carbonyl-containing compounds to their corresponding alcohols (PubMed: <a href="#">10510318</a> , PubMed: <a href="#">30538128</a> ). Displays enzymatic activity towards endogenous metabolites such as aromatic and aliphatic aldehydes, ketones, monosaccharides and bile acids, with a preference for negatively charged substrates, such as glucuronate and succinic semialdehyde (PubMed: <a href="#">10510318</a> , PubMed: <a href="#">30538128</a> ). Functions as a detoxifying enzyme by reducing a range of toxic aldehydes (By similarity). Reduces methylglyoxal and 3-deoxyglucosone, which are present at elevated levels under hyperglycemic conditions and are cytotoxic (By similarity). Involved also in the detoxification of lipid-derived aldehydes like acrolein (By similarity). Plays a role in the activation of procarcinogens, such as polycyclic aromatic hydrocarbon trans-dihydrodiols, and in the metabolism of various xenobiotics

and drugs, including the anthracyclines doxorubicin (DOX) and daunorubicin (DAUN) (PubMed:[11306097](#), PubMed:[18276838](#)). Also acts as an inhibitor of protein S-nitrosylation by mediating degradation of S-nitroso-coenzyme A (S-nitroso-CoA), a cofactor required to S-nitrosylate proteins (PubMed:[30538128](#)). S-nitroso-CoA reductase activity is involved in reprogramming intermediary metabolism in renal proximal tubules, notably by inhibiting protein S-nitrosylation of isoform 2 of PKM (PKM2) (By similarity). Also acts as a S-nitroso-glutathione reductase by catalyzing the NADPH-dependent reduction of S-nitrosoglutathione (PubMed:[31649033](#)). Displays no reductase activity towards retinoids (By similarity).

#### Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9JII6}. Apical cell membrane {ECO:0000250|UniProtKB:Q9JII6}

#### Tissue Location

Widely expressed. Highly expressed in kidney, salivary gland and liver. Detected in trachea, stomach, brain, lung, prostate, placenta, mammary gland, small intestine and lung


## Background

Catalyzes the NADPH-dependent reduction of a variety of aromatic and aliphatic aldehydes to their corresponding alcohols. Catalyzes the reduction of mevaldate to mevalonic acid and of glyceraldehyde to glycerol. Has broad substrate specificity. In vitro substrates include succinic semialdehyde, 4-nitrobenzaldehyde, 1,2-naphthoquinone, methylglyoxal, and D-glucuronic acid. Plays a role in the activation of procarcinogens, such as polycyclic aromatic hydrocarbon trans-dihydrodiols, and in the metabolism of various xenobiotics and drugs, including the anthracyclines doxorubicin (DOX) and daunorubicin (DAUN).

## References

Bohren K.M., et al. *J. Biol. Chem.* 264:9547-9551(1989).  
Fujii J., et al. *Cytogenet. Cell Genet.* 84:230-232(1999).  
Barski O.A., et al. *Genomics* 60:188-198(1999).  
Ota T., et al. *Nat. Genet.* 36:40-45(2004).  
Ebert L., et al. Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

## Images



Anti-AKR1A1 Antibody at 1:1000 dilution + human kidney lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 37 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.