

Cytochrome P450 2D6 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51914

Product Information

Application	WB
Primary Accession	P10635
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55769

Additional Information

Gene ID	1565
Other Names	Cytochrome P450 2D6, CYP2D6, Cytochrome P450-DB1, Debrisoquine 4-hydroxylase, CYP2D6, CYP2DL1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cytochrome P450 2D6. The exact sequence is proprietary.
Dilution	WB~~1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CYP2D6 {ECO:0000303 PubMed:21289075, ECO:0000312 HGNC:HGNC:2625}
Function	A cytochrome P450 monooxygenase involved in the metabolism of fatty acids, steroids and retinoids (PubMed:18698000, PubMed:19965576, PubMed:20972997, PubMed:21289075, PubMed:21576599). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:18698000, PubMed:19965576, PubMed:20972997, PubMed:21289075, PubMed:21576599). Catalyzes the epoxidation of double bonds of polyunsaturated fatty acids (PUFA) (PubMed:19965576, PubMed:20972997). Metabolizes endocannabinoid arachidonylethanolamide (anandamide) to 20-hydroxyeicosatetraenoic acid ethanolamide (20-HETE-EA) and 8,9-, 11,12-, and 14,15-epoxyeicosatrienoic acid ethanolamides (EpETRE-EAs), potentially modulating endocannabinoid system signaling

(PubMed:[18698000](#), PubMed:[21289075](#)). Catalyzes the hydroxylation of carbon-hydrogen bonds. Metabolizes cholesterol toward 25-hydroxycholesterol, a physiological regulator of cellular cholesterol homeostasis (PubMed:[21576599](#)). Catalyzes the oxidative transformations of all-trans retinol to all-trans retinal, a precursor for the active form all-trans-retinoic acid (PubMed:[10681376](#)). Also involved in the oxidative metabolism of drugs such as antiarrhythmics, adrenoceptor antagonists, and tricyclic antidepressants.

Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

Background

Responsible for the metabolism of many drugs and environmental chemicals that it oxidizes. It is involved in the metabolism of drugs such as antiarrhythmics, adrenoceptor antagonists, and tricyclic antidepressants.

References

- Gonzalez F.J.,et al.Genomics 2:174-179(1988).
Gonzalez F.J.,et al.Nature 331:442-446(1988).
Kimura S.,et al.Am. J. Hum. Genet. 45:889-904(1989).
Gaedigk A.,et al.Pharmacogenomics J. 5:173-182(2005).
Gaedigk A.,et al.Pharmacogenomics J. 5:276-276(2005).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.