

TLR2 Rabbit pAb

TLR2 Rabbit pAb Catalog # AP52045

Product Information

ApplicationWBPrimary Accession060603

Reactivity Human, Mouse

HostRabbitClonalityPolyclonalCalculated MW89838Physical StateLiquid

Immunogen KLH conjugated synthetic peptide derived from human TLR2

Epitope Specificity 701-784/784

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Membrane; Single-pass type I membrane protein.

SIMILARITY Belongs to the Toll-like receptor family.Contains 14 LRR (leucine-rich)

repeats. Contains 1 TIR domain.

SUBUNIT Interacts with LY96, TLR1 and TLR6 (via extracellular domain). Binds MYD88

(via TIR domain). Interacts with TICAM1. Ligand binding induces the formation

of a heterodimer with TLR1. Interacts with CNPY3.

Post-translational Glycosylation of Asn-442 is critical for secretion of the N-terminal ectodomain of TLR2.

ounications of the

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions The protein encoded by this gene is a member of the Toll-like receptor (TLR)

family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans

and share structural and functional similarities. They recognize

pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the

development of effective immunity. The various TLRs exhibit different patterns of expression. This gene is expressed most abundantly in peripheral blood leukocytes, and mediates host response to Gram-positive bacteria and

yeast via stimulation of NF-kappaB. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID 7097

Other Names Toll-like receptor 2, Toll/interleukin-1 receptor-like protein 4, CD282, TLR2

(HGNC:11848), TIL4

Target/Specificity Highly expressed in peripheral blood leukocytes, in particular in monocytes,

in bone marrow, lymph node and in spleen. Also detected in lung and in fetal

liver. Levels are low in other tissues.

Dilution WB=1:500-2000,Flow-Cyt=1 \(\text{Ig/Test} \)

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name TLR2 (<u>HGNC:11848</u>)

Synonyms TIL4

Function

Cooperates with LY96 to mediate the innate immune response to bacterial lipoproteins and other microbial cell wall components. Cooperates with TLR1 or TLR6 to mediate the innate immune response to bacterial lipoproteins or lipopeptides (PubMed: 17889651, PubMed: 21078852). Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. May also activate immune cells and promote apoptosis in response to the lipid moiety of lipoproteins (PubMed: 10426995, PubMed: 10426996). Recognizes mycoplasmal macrophage-activating lipopeptide-2kD (MALP-2), soluble tuberculosis factor (STF), phenol-soluble modulin (PSM) and B.burgdorferi outer surface protein A lipoprotein (OspA-L) cooperatively with TLR6 (PubMed: 11441107). Stimulation of monocytes in vitro with M.tuberculosis PstS1 induces p38 MAPK and ERK1/2 activation primarily via this receptor, but also partially via TLR4 (PubMed: 16622205). MAPK activation in response to bacterial peptidoglycan also occurs via this receptor (PubMed: 16622205). Acts as a receptor for M.tuberculosis lipoproteins LprA, LprG, LpqH and PstS1, some lipoproteins are dependent on other coreceptors (TLR1, CD14 and/or CD36); the lipoproteins act as agonists to modulate antigen presenting cell functions in response to the pathogen (PubMed:19362712). M.tuberculosis HSP70 (dnaK) but not HSP65 (groEL-2) acts via this protein to stimulate NF-kappa-B expression (PubMed: 15809303). Recognizes M.tuberculosis major T-antigen EsxA (ESAT-6) which inhibits downstream MYD88-dependent signaling (shown in mouse) (By similarity). Forms activation clusters composed of several receptors depending on the ligand, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway. Forms the cluster TLR2:TLR6:CD14:CD36 in response to diacylated lipopeptides and TLR2:TLR1:CD14 in response to triacylated lipopeptides (PubMed:<u>16880211</u>). Required for normal uptake of M.tuberculosis, a process that is inhibited by M.tuberculosis LppM (By similarity).

Cellular Location

Membrane {ECO:0000250 | UniProtKB:Q9QUN7}; Single- pass type I membrane protein. Cytoplasmic vesicle, phagosome membrane {ECO:0000250 | UniProtKB:Q9QUN7}; Single-pass type I membrane protein. Membrane raft. Note=Does not reside in lipid rafts before stimulation but accumulates increasingly in the raft upon the presence of the microbial ligand. In response to diacylated lipoproteins, TLR2:TLR6 heterodimers are recruited in lipid rafts, this recruitment determines the intracellular targeting to the Golgi apparatus. Triacylated lipoproteins induce the same mechanism for TLR2:TLR1 heterodimers.

Tissue Location

Highly expressed in peripheral blood leukocytes, in particular in monocytes, in bone marrow, lymph node and in spleen. Also detected in lung and in fetal liver. Levels are low in other tissues


Background

The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This gene is expressed most abundantly in peripheral blood leukocytes, and mediates host response to Gram-positive bacteria and yeast via stimulation of NF-kappaB. [provided by RefSeq, Jul 2008].


References

Chaudhary P.M., et al. Blood 91:4020-4027(1998). Rock F.L., et al. Proc. Natl. Acad. Sci. U.S.A. 95:588-593(1998). Yang R.-B., et al. Nature 395:284-288(1998). Nakajima T., et al. Immunogenetics 60:727-735(2008). Georgel P., et al. PLoS ONE 4:E7803-E7803(2009).

Images



The THP-1 (H) cells were fixed with 4% PFA (10 min at r.t.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then were incubated in 5%BSA to block non-specific protein-protein interactions (30 min at r.t.). Primary Antibody (green): Rabbit Anti-TLR2 antibody (AP52045): 1 μ g/10^6 cells; Secondary Antibody (white blue): Goat anti-Rabbit IgG-BF488 (AP52045-BF488): 1 μ g/test. Blank control (black): PBS. Acquisition of 20,000 events was performed.



25 ug total protein per lane of various lysates (see on figure) probed with TLR2 polyclonal antibody, unconjugated (AP52045) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

Citations

• Deletion of Thioredoxin-interacting protein ameliorates high fat diet-induced non-alcoholic steatohepatitis through modulation of Toll-like receptor 2-NLRP3-inflammasome axis: Histological and immunohistochemical study.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.