

ROCK2 Rabbit pAb

ROCK2 Rabbit pAb
Catalog # AP52048

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q75116
Reactivity	Human, Mouse
Predicted	Rat, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	160900
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ROCK2
Epitope Specificity	1001-1300/1388
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Cell membrane. Cytoplasmic, and associated with actin microfilaments and the plasma membrane.
SIMILARITY	Belongs to the protein kinase superfamily. AGC Ser/Thrprotein kinase family. Contains 1 AGC-kinase C-terminal domain. Contains 1 PH domain.
SUBUNIT	Homodimer. Interacts with IRS1, RHOB and RHOC. Interacts with RHOA (activated by GTP), PPP1R12A, CHORDC1, SORL1, EP300 and BRCA2. Interacts with NPM1 and this interaction enhances its activity. Interacts with RAF1.
Post-translational modifications	Phosphorylated upon DNA damage, probably by ATM or ATR. Phosphorylation at Tyr-722 reduces its binding to RHOA and is crucial for focal adhesion dynamics. Dephosphorylation by PTPN11 stimulates its RHOA binding activity.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is a serine/threonine kinase that regulates cytokinesis, smooth muscle contraction, the formation of actin stress fibers and focal adhesions, and the activation of the c-fos serum response element. This protein, which is an isozyme of ROCK1 is a target for the small GTPase Rho. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID	9475
Other Names	Rho-associated protein kinase 2, 2.7.11.1, Rho kinase 2, Rho-associated, coiled-coil-containing protein kinase 2, Rho-associated, coiled-coil-containing protein kinase II, ROCK-II, p164 ROCK-2, ROCK2, KIAA0619
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	ROCK2
Synonyms	KIAA0619
Function	Protein kinase which is a key regulator of actin cytoskeleton and cell polarity. Involved in regulation of smooth muscle contraction, actin cytoskeleton organization, stress fiber and focal adhesion formation, neurite retraction, cell adhesion and motility via phosphorylation of ADD1, BRCA2, CNN1, EZR, DPYSL2, EP300, MSN, MYL9/MLC2, NPM1, RDX, PPP1R12A and VIM. Phosphorylates SORL1 and IRF4. Acts as a negative regulator of VEGF-induced angiogenic endothelial cell activation. Positively regulates the activation of p42/MAPK1- p44/MAPK3 and of p90RSK/RPS6KA1 during myogenic differentiation. Plays an important role in the timely initiation of centrosome duplication. Inhibits keratinocyte terminal differentiation. May regulate closure of the eyelids and ventral body wall through organization of actomyosin bundles. Plays a critical role in the regulation of spine and synaptic properties in the hippocampus. Plays an important role in generating the circadian rhythm of the aortic myofilament Ca(2+) sensitivity and vascular contractility by modulating the myosin light chain phosphorylation.
Cellular Location	Cytoplasm. Cell membrane; Peripheral membrane protein. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Cytoplasmic, and associated with actin microfilaments and the plasma membrane.
Tissue Location	Expressed in the brain (at protein level).

Background


The protein encoded by this gene is a serine/threonine kinase that regulates cytokinesis, smooth muscle contraction, the formation of actin stress fibers and focal adhesions, and the activation of the c-fos serum response element. This protein, which is an isozyme of ROCK1 is a target for the small GTPase Rho. [provided by RefSeq, Jul 2008].

References


Takahashi N.,et al.Genomics 55:235-237(1999).
Ishikawa K.,et al.DNA Res. 5:169-176(1998).
Hillier L.W.,et al.Nature 434:724-731(2005).
Kawano Y.,et al.J. Cell Biol. 147:1023-1038(1999).
Sebbagh M.,et al.J. Exp. Med. 201:465-471(2005).

Images

Sample:
Lane 1: Mouse Heart Lysates
Lane 2: Mouse Skeletal muscle Lysates



Primary: Anti-ROCK2 (AP52048) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 161kDa
 Observed band size: 180kDa



Tissue/cell: human gastric carcinoma; 4%
 Paraformaldehyde-fixed and paraffin-embedded;
 Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling
 bathing for 15min; Block endogenous peroxidase by 3%
 Hydrogen peroxide for 30min; Blocking buffer (normal
 goat serum,C-0005) at 37°C for 20 min;
 Incubation: Anti-ROCK2 Polyclonal Antibody,
 Unconjugated(AP52048) 1:200, overnight at 4°C, followed
 by conjugation to the secondary antibody(SP-0023) and
 DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.