

Pyruvate Dehydrogenase E2 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AP52757

Product Information

Application	WB, ICC, IP
Primary Accession	P10515
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	68997

Additional Information

Gene ID	1737
Other Names	70 kDa mitochondrial autoantigen of primary biliary cirrhosis;anti DLAT; Dihydrolipoamide acetyltransferase component of pyruvate dehydrogenase complex; Dihydrolipoamide;Dihydrolipoamide S Acetyltransferase;Dihydrolipoamide S-acetyltransferase (E2 component of pyruvate dehydrogenase complex);Dihydrolipoamide S-Acetyltransferase;Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex;dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex mitochondrial;DLAT;DLAT;DLTA;E2;E2 component of pyruvate dehydrogenase complex;EC 2.3.1.12;M2 antigen complex 70 kDa subunit;M2 Antigen Complex 70kD Subunit;mitochondrial;ODP2_HUMAN;PBC;PDC E2;PDC-E2;PDCE2;Pyruvate dehydrogenase complex component E2;Pyruvate dehydrogenase complex E2 subunit;S acetyltransferase component of pyruvate dehydrogenase complex.
Dilution	WB~~1:1000 ICC~~1:300 IP~~1:500
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	DLAT (HGNC:2896)
Synonyms	DLTA The pyruvate dehydrogenase (PDH) complex, catalyzes the overall

Function conversion of pyruvate to acetyl-CoA and CO₂, and thereby links cytoplasmic glycolysis and the mitochondrial tricarboxylic acid (TCA) cycle (Probable). It contains multiple copies of three enzymatic components: pyruvate dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and dihydrolipoamide dehydrogenase (E3); (Probable). Within this complex, the catalytic function of this enzyme is to accept, and to transfer to coenzyme A, acetyl groups from acetyl- lipoyl moiety generated by the pyruvate dehydrogenase, leading to acetyl-CoA formation (Probable).

Cellular Location Mitochondrion matrix {ECO:0000250|UniProtKB:P08461}


Background

The pyruvate dehydrogenase complex catalyzes the overall conversion of pyruvate to acetyl-CoA and CO₂, and thereby links the glycolytic pathway to the tricarboxylic cycle.


References

Totoki Y., et al. Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Taylor T.D., et al. Nature 440:497-500(2006).
Coppel R.L., et al. Proc. Natl. Acad. Sci. U.S.A. 85:7317-7321(1988).
Thekkumkara T.J., et al. FEBS Lett. 240:45-48(1988).
Hiromasa Y., et al. J. Biol. Chem. 279:6921-6933(2004).

Images




Western blot detection of Pyruvate Dehydrogenase E2 in Jurkat, A549, U251, F9, Lncap and HeLa cell lysates using Pyruvate Dehydrogenase E2 mouse mAb (1:1000 diluted). Predicted band size: 69KDa. Observed band size: 69KDa.



Immunocytochemistry stain of HeLa using Pyruvate Dehydrogenase E2 mouse mAb (1:300).

Immunoprecipitation analysis of HeLa cell lysates using Pyruvate Dehydrogenase E2 mouse mAb.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.