

HDAC3 Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52764

Product Information

Application	WB
Primary Accession	O15379
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	48848

Additional Information

Gene ID	8841
Other Names	HD 3;HD3;HDAC 3;HDAC3;HDAC3_HUMAN;Histone deacetylase 3 (HD3) (RPD3-2);histone deacetylase 3;RPD 3;RPD3;RPD3-2;SMAP 45;SMAP45.
Dilution	WB~1:1000
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	HDAC3
Function	Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4), and some other non-histone substrates (PubMed: 21030595 , PubMed: 21444723 , PubMed: 23911289 , PubMed: 25301942 , PubMed: 28167758 , PubMed: 28497810 , PubMed: 32404892 , PubMed: 22230954). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed: 23911289). Histone deacetylases act via the formation of large multiprotein complexes, such as N-Cor repressor complex, which activate the histone deacetylase activity (PubMed: 23911289 , PubMed: 22230954). Participates in the BCL6 transcriptional repressor activity by deacetylating the H3 'Lys-27' (H3K27) on enhancer elements, antagonizing EP300 acetyltransferase activity and repressing proximal gene expression (PubMed: 23911289). Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation (By similarity).

Contributes, together with XBP1 isoform 1, to the activation of NFE2L2-mediated HMOX1 transcription factor gene expression in a PI(3)K/mTORC2/Akt-dependent signaling pathway leading to endothelial cell (EC) survival under disturbed flow/oxidative stress (PubMed:[25190803](#)). Regulates both the transcriptional activation and repression phases of the circadian clock in a deacetylase activity-independent manner (By similarity). During the activation phase, promotes the accumulation of ubiquitinated BMAL1 at the E-boxes and during the repression phase, blocks FBXL3-mediated CRY1/2 ubiquitination and promotes the interaction of CRY1 and BMAL1 (By similarity). The NCOR1-HDAC3 complex regulates the circadian expression of the core clock gene BMAL1 and the genes involved in lipid metabolism in the liver (By similarity). Also functions as a deacetylase for non-histone targets, such as KAT5, MEF2D, MAPK14, RARA and STAT3 (PubMed:[15653507](#), PubMed:[21030595](#), PubMed:[21444723](#), PubMed:[25301942](#), PubMed:[28167758](#)). Serves as a corepressor of RARA, mediating its deacetylation and repression, leading to inhibition of RARE DNA element binding (PubMed:[28167758](#)). In association with RARA, plays a role in the repression of microRNA-10a and thereby in the inflammatory response (PubMed:[28167758](#)). In addition to protein deacetylase activity, also acts as a protein-lysine deacetylase by recognizing other acyl groups: catalyzes removal of (2E)-butenoyl (crotonyl), lactoyl (lactyl), 2-hydroxyisobutanoyl (2-hydroxyisobutyryl) and isonicotinyl acyl groups from lysine residues, leading to protein deacetylation, delactylation, de-2-hydroxyisobutyrylation and deisonicotinylation, respectively (PubMed:[28497810](#), PubMed:[29192674](#), PubMed:[34608293](#), PubMed:[34545082](#), PubMed:[35044827](#)). Catalyzes deacetylation of MAPRE1/EB1 (PubMed:[34608293](#)). Mediates delactylation NBN/NBS1, thereby inhibiting DNA double-strand breaks (DSBs) via homologous recombination (HR) (PubMed:[38961290](#)).

Cellular Location	Nucleus. Chromosome. Cytoplasm. Cytoplasm, cytosol. Note=Colocalizes with XBP1 and AKT1 in the cytoplasm (PubMed:25190803). Predominantly expressed in the nucleus in the presence of CCAR2 (PubMed:21030595)
Tissue Location	Widely expressed..


Background

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4), and some other non-histone substrates. Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Participates in the BCL6 transcriptional repressor activity by deacetylating the H3 'Lys- 27' (H3K27) on enhancer elements, antagonizing EP300 acetyltransferase activity and repressing proximal gene expression. Probably participates in the regulation of transcription through its binding to the zinc-finger transcription factor YY1; increases YY1 repression activity. Required to repress transcription of the POU1F1 transcription factor. Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation (PubMed:[21444723](#), PubMed:[23911289](#)). Contributes, together with XBP1 isoform 1, to the activation of NFE2L2-mediated HMOX1 transcription factor gene expression in a PI(3)K/mTORC2/Akt-dependent signaling pathway leading to endothelial cell (EC) survival under disturbed flow/oxidative stress (PubMed:[25190803](#)).

References

Dangond F.,et al.Biochem. Biophys. Res. Commun. 242:648-652(1998).
Yang W.-M.,et al.J. Biol. Chem. 272:28001-28007(1997).
Emiliani S.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:2795-2800(1998).
Mahlknecht U.,et al.Genomics 56:197-202(1999).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Images



Western blot detection of HDAC3 in THP-1,LNCAP,MCF7,PC12,293,Jurkat,3T3 and Hela cell lysates using HDAC3 mouse mAb (1:1000 diluted).Predicted band size:49KDa.Observed band size:49KDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.