

Anti-MC4 Receptor Antibody

Catalog # AP53688

Product Information

| | |
|--------------------------|------------------------|
| Application | WB |
| Primary Accession | P32245 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 36943 |

Additional Information

| | |
|---------------------------|---|
| Gene ID | 4160 |
| Other Names | Melanocortin receptor 4; MC4-R |
| Target/Specificity | Recognizes endogenous levels of MC4 Receptor protein. |
| Dilution | WB~~1/500 - 1/1000 |
| Format | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide. |
| Storage | Store at -20 °C. Stable for 12 months from date of receipt |

Protein Information

| | |
|-----------------|--|
| Name | MC4R (HGNC:6932) |
| Function | G protein-coupled receptor that binds melanocyte-stimulating hormones (alpha- and beta-MSH) and corticotropin/ACTH, which are peptide products of the POMC precursor (PubMed: 12646665 , PubMed: 14764818 , PubMed: 25163632 , PubMed: 32327598 , PubMed: 33858992 , PubMed: 8392067). Functions as a central component of the leptin-melanocortin pathway, which is essential for maintaining energy homeostasis (PubMed: 32327598 , PubMed: 33858992). Upon activation, couples to G(s) protein, stimulating adenylate cyclase and the cAMP- dependent signaling pathway, which promotes anorexogenic signaling in the hypothalamus and contributes to a negative energy balance (PubMed: 12588803 , PubMed: 14764818 , PubMed: 25163632 , PubMed: 33858992). Regulates food intake: activation by agonists suppresses appetite, whereas the antagonist Agouti-related protein/AGRP precludes agonist- induced signaling, thereby stimulating appetite (PubMed: 9311920). Modulates the firing activity of neurons in paraventricular nucleus (PVN) of the hypothalamus via alpha-MSH and AGRP regulation of inwardly rectifying potassium channel KCNJ13 closure, independently of G(s) signaling (PubMed: 32327598). In the PVN, also interacts |

with opsin 3/OPN3, which couples to G(i/o) proteins to inhibit MC4R-mediated cAMP signaling, thereby promoting food intake (PubMed:[39951488](#)). In intestinal epithelial cells, contributes to inhibition of hepatic glucose production via nesfatin-1/NUCB2, leading to increased cAMP levels and glucagon-like peptide 1 (GLP-1) secretion (PubMed:[39562740](#)). Interaction with MGRN1 displaces the G(s) protein, further decreasing MC4R signaling activity (PubMed:[19737927](#)). Also activated by gamma-MSH, though with low potency (PubMed:[8392067](#)).


Cellular Location Cell membrane; Multi-pass membrane protein

Tissue Location Brain, placental, and gut tissues.


Background

Rabbit polyclonal antibody to MC4 Receptor

Images



Western blot analysis of MC4 Receptor expression in mouse brain (A) whole cell lysates.



Immunohistochemical analysis of MC4 Receptor staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.