

# Anti-CTH Antibody

Catalog # AP53807

## Product Information

---

Application	WB
Primary Accession	<a href="#">P32929</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44508

## Additional Information

---

Gene ID	1491
Other Names	Cystathionine gamma-lyase; Cysteine-protein sulfhydrase; Gamma-cystathionase
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CTH. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/1000
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	CTH
Function	Catalyzes the last step in the trans-sulfuration pathway from L-methionine to L-cysteine in a pyridoxal-5'-phosphate (PLP)-dependent manner, which consists on cleaving the L,L-cystathionine molecule into L-cysteine, ammonia and 2-oxobutanoate (PubMed: <a href="#">10212249</a> , PubMed: <a href="#">18476726</a> , PubMed: <a href="#">19261609</a> , PubMed: <a href="#">19961860</a> ). Part of the L- cysteine derived from the trans-sulfuration pathway is utilized for biosynthesis of the ubiquitous antioxidant glutathione (PubMed: <a href="#">18476726</a> ). Besides its role in the conversion of L- cystathionine into L-cysteine, it utilizes L-cysteine and L- homocysteine as substrates (at much lower rates than L,L-cystathionine) to produce the endogenous gaseous signaling molecule hydrogen sulfide (H2S) (PubMed: <a href="#">10212249</a> , PubMed: <a href="#">19019829</a> , PubMed: <a href="#">19261609</a> , PubMed: <a href="#">19961860</a> ). In vitro, it converts two L-cysteine molecules into lanthionine and H2S, also two L-homocysteine molecules to homolanthionine and H2S, which can be particularly relevant under conditions of severe hyperhomocysteinemia (which is a risk factor for cardiovascular disease,

diabetes, and Alzheimer's disease) (PubMed:[19261609](#)). Lanthionine and homolanthionine are structural homologs of L,L-cystathionine that differ by the absence or presence of an extra methylene group, respectively (PubMed:[19261609](#)). Acts as a cysteine-protein sulfhydrase by mediating sulfhydration of target proteins: sulfhydration consists of converting -SH groups into -SSH on specific cysteine residues of target proteins such as GAPDH, PTPN1 and NF-kappa-B subunit RELA, thereby regulating their function (PubMed:[22169477](#)). By generating the gasotransmitter H<sub>2</sub>S, it participates in a number of physiological processes such as vasodilation, bone protection, and inflammation (Probable) (PubMed:[29254196](#)). Plays an essential role in myogenesis by contributing to the biogenesis of H<sub>2</sub>S in skeletal muscle tissue (By similarity). Can also accept homoserine as substrate (By similarity). Catalyzes the elimination of selenocystathionine (which can be derived from the diet) to yield selenocysteine, ammonia and 2-oxobutanoate (By similarity).

**Cellular Location**

Cytoplasm.

**Tissue Location**

Highly expressed in liver (PubMed:10727430, PubMed:20305127). Also in muscle and lower expression in most tissues except heart, pituitary gland, spleen, thymus, and vascular tissue, where it is hardly detected (PubMed:20305127)


## Background

---

Rabbit polyclonal antibody to CTH

## Images

---



Western blot analysis of CTH expression in HEK293T (A), Raw264.7 (B), PC12 (C) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.