

Anti-Girdin Antibody

Catalog # AP54006

Product Information

Application	WB
Primary Accession	Q3V6T2
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	216042

Additional Information

Gene ID	55704
Other Names	APE; GRDN; KIAA1212; Girdin; Akt phosphorylation enhancer; APE; Coiled-coil domain-containing protein 88A; G alpha-interacting vesicle-associated protein; GIV; Girders of actin filament; Hook-related protein 1; HkRP1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Girdin. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/1000
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CCDC88A
Function	Bifunctional modulator of guanine nucleotide-binding proteins (G proteins) (PubMed: 19211784 , PubMed: 27621449). Acts as a non-receptor guanine nucleotide exchange factor which binds to and activates guanine nucleotide-binding protein G(i) alpha subunits (PubMed: 19211784 , PubMed: 21954290 , PubMed: 23509302 , PubMed: 25187647). Also acts as a guanine nucleotide dissociation inhibitor for guanine nucleotide-binding protein G(s) subunit alpha GNAS (PubMed: 27621449). Essential for cell migration (PubMed: 16139227 , PubMed: 19211784 , PubMed: 20462955 , PubMed: 21954290). Interacts in complex with G(i) alpha subunits with the EGFR receptor, retaining EGFR at the cell membrane following ligand stimulation and promoting EGFR signaling which triggers cell migration (PubMed: 20462955). Binding to Gi-alpha subunits displaces the beta and gamma subunits from the heterotrimeric G-protein complex which enhances phosphoinositide 3-kinase (PI3K)-dependent phosphorylation and kinase

activity of AKT1/PKB (PubMed:[19211784](#)). Phosphorylation of AKT1/PKB induces the phosphorylation of downstream effectors GSK3 and FOXO1/FKHR, and regulates DNA replication and cell proliferation (By similarity). Binds in its tyrosine-phosphorylated form to the phosphatidylinositol 3-kinase (PI3K) regulatory subunit PIK3R1 which enables recruitment of PIK3R1 to the EGFR receptor, enhancing PI3K activity and cell migration (PubMed:[21954290](#)). Plays a role as a key modulator of the AKT-mTOR signaling pathway, controlling the tempo of the process of newborn neuron integration during adult neurogenesis, including correct neuron positioning, dendritic development and synapse formation (By similarity). Inhibition of G(s) subunit alpha GNAS leads to reduced cellular levels of cAMP and suppression of cell proliferation (PubMed:[27621449](#)). Essential for the integrity of the actin cytoskeleton (PubMed:[16139227](#), PubMed:[19211784](#)). Required for formation of actin stress fibers and lamellipodia (PubMed:[15882442](#)). May be involved in membrane sorting in the early endosome (PubMed:[15882442](#)). Plays a role in ciliogenesis and cilium morphology and positioning and this may partly be through regulation of the localization of scaffolding protein CROCC/Rootletin (PubMed:[27623382](#)).

Cellular Location

Cell membrane; Peripheral membrane protein. Cytoplasm, cytosol. Cytoplasmic vesicle. Cell projection, lamellipodium. Cytoplasm, cytoskeleton, cilium basal body. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Note=Localizes to the cytosol in unstimulated cells while EGF stimulation promotes membrane localization and guanine nucleotide exchange factor activity (PubMed:[27864364](#)) Localizes to the cell membrane through interaction with phosphoinositides (PubMed:[15882442](#), PubMed:[16139227](#))


Tissue Location

Expressed ubiquitously.

Background


Rabbit polyclonal antibody to Girdin

Images



Western blot analysis of Girdin expression in K562 (A), CT26 (B), rat testis (C) whole cell lysates.

Immunohistochemical analysis of Girdin staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.