

Anti-OCT1 Antibody

Catalog # AP54135

Product Information

Application WB
Primary Accession P14859
Other Accession P09086
Reactivity Human
Host Mouse
Clonality Monoclonal
Calculated MW 76472

Additional Information

Gene ID 5451

Other Names OCT1; OTF1; POU domain class 2 transcription factor 1; NF-A1;

Octamer-binding protein 1; Oct-1; Octamer-binding transcription factor 1;

OTF-1

Target/Specificity KLH-conjugated synthetic peptide encompassing a sequence of human OCT1.

The exact sequence is proprietary.

Dilution WB~~1/500 - 1/2000

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name POU2F1

Synonyms OCT1, OTF1

Function Transcription factor that binds to the octamer motif (5'- ATTTGCAT-3') and

activates the promoters of the genes for some small nuclear RNAs (snRNA) and of genes such as those for histone H2B and immunoglobulins. Modulates

transcription transactivation by NR3C1, AR and PGR.


Cellular Location Nucleus.

Tissue Location Ubiquitous. Isoform 2 is lymphocyte-specific.


Background

Mouse monoclonal antibody to OCT1

Images



Western blot analysis of 42278 expression in Hela (A), Jurkat (B), HepG2 (C) whole cell lysates.



Immunohistochemical analysis of 42278 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.