

# EXOC3 Rabbit pAb

EXOC3 Rabbit pAb  
Catalog # AP54486

## Product Information

---

<b>Primary Accession</b>	<a href="#">O60645</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Mouse, Rat, Dog, Rabbit, Sheep
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	85567
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated synthetic peptide derived from human EXOC3
<b>Epitope Specificity</b>	714-756/756
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Cytoplasmic
<b>SIMILARITY</b>	Belongs to the SEC6 family.
<b>SUBUNIT</b>	The exocyst complex is composed of EXOC1, EXOC2, EXOC3, EXOC4, EXOC5, EXOC6, EXOC7 and EXOC8. Interacts with EXOC3L1 (By similarity). Interacts with BIRC6/bruce. Interacts with MYRIP
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	The protein encoded by this gene is a component of the exocyst complex, a multiple protein complex essential for targeting exocytic vesicles to specific docking sites on the plasma membrane. Though best characterized in yeast, the component proteins and functions of exocyst complex have been demonstrated to be highly conserved in higher eukaryotes. At least eight components of the exocyst complex, including this protein, are found to interact with the actin cytoskeletal remodeling and vesicle transport machinery. The complex is also essential for the biogenesis of epithelial cell surface polarity.

## Additional Information

---

<b>Gene ID</b>	11336
<b>Other Names</b>	Exocyst complex component 3, Exocyst complex component Sec6, EXOC3, SEC6, SEC6L1
<b>Dilution</b>	Flow-Cyt=2ug/Test
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


## Protein Information

<b>Name</b>	EXOC3
<b>Synonyms</b>	SEC6, SEC6L1
<b>Function</b>	Component of the exocyst complex involved in the docking of exocytic vesicles with fusion sites on the plasma membrane.
<b>Cellular Location</b>	Cytoplasm {ECO:0000250 UniProtKB:O54921}. Cytoplasm, perinuclear region {ECO:0000250 UniProtKB:O54921}. Cell projection, growth cone {ECO:0000250 UniProtKB:O54921}. Midbody. Golgi apparatus. Cell projection, neuron projection {ECO:0000250 UniProtKB:Q62825}. Note=Perinuclear in undifferentiated cells. Redistributes to growing neurites and growth cones during neuronal differentiation (By similarity). During mitosis, early recruitment to the midbody requires RALA, but not RALB, and EXOC2. In late stages of cytokinesis, localization to the midbody is RALB- dependent (PubMed:18756269). {ECO:0000250 UniProtKB:O54921, ECO:0000269 PubMed:18756269}
<b>Tissue Location</b>	Expressed in epididymis (at protein level).

## Background


The protein encoded by this gene is a component of the exocyst complex, a multiple protein complex essential for targeting exocytic vesicles to specific docking sites on the plasma membrane. Though best characterized in yeast, the component proteins and functions of exocyst complex have been demonstrated to be highly conserved in higher eukaryotes. At least eight components of the exocyst complex, including this protein, are found to interact with the actin cytoskeletal remodeling and vesicle transport machinery. The complex is also essential for the biogenesis of epithelial cell surface polarity.

## Images



Blank control:HeLa.  
Primary Antibody (green line): Rabbit Anti-rSec6 antibody (AP54486)  
Dilution: 2  $\mu\text{g}$  / $10^6$  cells;  
Isotype Control Antibody (orange line): Rabbit IgG .  
Secondary Antibody : Goat anti-rabbit IgG-PE  
Dilution: 1  $\mu\text{g}$  /test.  
Protocol  
The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Blank control:HeLa.  
Primary Antibody (green line): Rabbit Anti-rSec6 antibody (AP54486)  
Dilution: 2  $\mu\text{g}$  / $10^6$  cells;  
Isotype Control Antibody (orange line): Rabbit IgG .  
Secondary Antibody : Goat anti-rabbit IgG-PE



Dilution: 1 µg /test.

Protocol

The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.