

5HT7 Receptor Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54714

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	P34969
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53555
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human 5HT7 Receptor/SR-7
Epitope Specificity	51-150/479
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane; Multi-pass membrane protein.
SIMILARITY	Belongs to the G-protein coupled receptor 1 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The neurotransmitter, serotonin, is thought to play a role in various cognitive and behavioral functions. The serotonin receptor encoded by this gene belongs to the superfamily of G protein-coupled receptors and the gene is a candidate locus for involvement in autistic disorder and other neuropsychiatric disorders. Three splice variants have been identified which encode proteins that differ in the length of their carboxy terminal ends. [provided by RefSeq, Jul 2008]

Additional Information


Gene ID	3363
Other Names	5-hydroxytryptamine receptor 7, 5-HT-7, 5-HT7, 5-HT-X, Serotonin receptor 7, HTR7
Target/Specificity	Isoform A is the predominant isoform in spleen, caudate and hippocampus. Isoform B is expressed at lower levels. Isoform D is a minor isoform in term of expression.
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	HTR7 {ECO:0000303 Ref.3, ECO:0000312 HGNC:HGNC:5302}
Function	G-protein coupled receptor for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone and a mitogen (PubMed: 35714614 , PubMed: 8226867). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors (PubMed: 35714614 , PubMed: 8226867). HTR7 is coupled to G(s) G alpha proteins and mediates activation of adenylate cyclase activity (PubMed: 35714614).
Cellular Location	Cell membrane; Multi-pass membrane protein
Tissue Location	[Isoform A]: Predominant isoform in spleen, caudate and hippocampus. [Isoform D]: Minor isoform in terms of expression.

Images



Sample:
Hippocampus (Mouse) Lysate at 40 ug
Hippocampus(Rat) Lysate at 40 ug
Primary: Anti- 5HT7 Receptor (AP54714) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 54 kD
Observed band size: 54 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.