

Actin Regulatory Protein CAPG Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54856

Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession P40121

Reactivity Rat, Dog, Bovine

HostRabbitClonalityPolyclonalCalculated MW38499

Additional Information

Gene ID 822

Other Names Macrophage-capping protein, Actin regulatory protein CAP-G, CAPG, AFCP,

MCP

Dilution IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-

10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name CAPG

Synonyms AFCP, MCP

Function Calcium-sensitive protein which reversibly blocks the barbed ends of actin

filaments but does not sever preformed actin filaments. May play an important role in macrophage function. May play a role in regulating cytoplasmic and/or nuclear structures through potential interactions with

actin. May bind DNA.

Cellular Location Nucleus. Cytoplasm Melanosome. Cell projection, lamellipodium

{ECO:0000250|UniProtKB:P24452}. Cell projection, ruffle

{ECO:0000250 | UniProtKB:P24452}. Note=In macrophages, may be

predominantly cytoplasmic. Nuclear localization was observed in fibroblasts. In macrophages, present at the membrane-cytoplasm interface. In activated

macrophages, concentrated in the ruffles of the leading lamellipodia.

{ECO:0000250 | UniProtKB:P24452}

Tissue Location

Macrophages and macrophage-like cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.