

HOXD8 Rabbit pAb

HOXD8 Rabbit pAb
Catalog # AP55215

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P13378
Reactivity	Human, Mouse
Predicted	Rat, Pig, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31911
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HOXD8
Epitope Specificity	151-250/290
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus
SIMILARITY	Belongs to the Antp homeobox family. Contains 1 homeobox DNA-binding domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The Hox proteins are a family of transcription factors that play a role in development and cellular differentiation by regulating downstream target genes. Specifically, the Hox proteins direct DNA-protein and protein-protein interactions that assist in determining the morphologic features associated with the anterior-posterior body axis. Hox proteins are involved in controlling axial patterning, leukemias and hereditary malformations. HoxD8 (homeobox D8), also known as HOX4E, is a 290 amino acid protein that localizes to the nucleus and contains one homeobox DNA-binding domain. One of several members of the homeobox superfamily, HoxD8 functions as a sequence-specific transcription factor that is important for the correct positioning of developing limb buds on the anterior-posterior axis.

Additional Information

Gene ID	3234
Other Names	Homeobox protein Hox-D8, Homeobox protein Hox-4E, Homeobox protein Hox-5.4, HOXD8, HOX4E
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


Protein Information

Name	HOXD8
Synonyms	HOX4E
Function	Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis.
Cellular Location	Nucleus.


Background

The Hox proteins are a family of transcription factors that play a role in development and cellular differentiation by regulating downstream target genes. Specifically, the Hox proteins direct DNA-protein and protein-protein interactions that assist in determining the morphologic features associated with the anterior-posterior body axis. Hox proteins are involved in controlling axial patterning, leukemias and hereditary malformations. HoxD8 (homeobox D8), also known as HOX4E, is a 290 amino acid protein that localizes to the nucleus and contains one homeobox DNA-binding domain. One of several members of the homeobox superfamily, HoxD8 functions as a sequence-specific transcription factor that is important for the correct positioning of developing limb buds on the anterior-posterior axis.

Images



Protein: muscle(mouse) lysate at 40ug;
Primary: rabbit Anti-HOXD8 (AP55215) at 1:300;
Secondary: HRP conjugated Goat-Anti-rabbit
IgG(AP55215-HRP) at 1: 5000;
Predicted band size: 32 kD
Observed band size: 34 kD



Tissue/cell: Human colon cancer; 4%
Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling
bathing for 15min; Block endogenous peroxidase by 3%
Hydrogen peroxide for 30min; Blocking buffer (normal
goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-HOXC8 Polyclonal Antibody,
Unconjugated(AP55215) 1:200, overnight at 4°C, followed
by conjugation to the secondary antibody(SP-0023) and
DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.