

EGR4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55609

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q05215
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61623
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human EGR4
Epitope Specificity	501-589/589
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nuclear
SIMILARITY	Belongs to the EGR C2H2-type zinc-finger protein family.Contains 3 C2H2-type zinc fingers.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Egr-1, Egr-2, Egr-3 and Egr-4 are nuclear transcription factors belonging to the Egr C2H2-type zinc-finger protein family and containing three C2H2-type zinc fingers. As immediate early proteins, Egr transcription factors are rapidly induced by diverse extracellular stimuli. They are subject to tight differential control through diverse mechanisms at several levels of regulation: transcriptional; translational and posttranslational (including glycosylation, phosphorylation and redox) mechanisms; and protein-protein interaction. Egr-1 binds to the DNA sequence 5'-CGCCCCGC-3' (Egr-site), thereby activating transcription of target genes whose products are required for mitogenesis and differentiation. Egr-2 binds specific DNA sites located in the promoter region of HoxA4. Egr-2 defects cause congenital hypomyelination neuropathy (also designated Charcot-Marie-Tooth disease) and Dejerine-Sottas neuropathology (also designated hereditary motor and sensory neuropathy III). Egr-3 is involved in muscle spindle development and is expressed in T cells 20 minutes following activation. Egr-4 binds to the Egr consensus motif GCGTGGGCG, functions as a transcriptional repressor, and displays autoregulatory activities, downregulating its own gene promoter in a dose dependent manner.

Additional Information

Gene ID	1961
Other Names	Early growth response protein 4, EGR-4, AT133, EGR4

Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	EGR4
Function	Transcriptional regulator. Recognizes and binds to the DNA sequence 5'-GCGGGGCG-3' (GSG). Activates the transcription of target genes whose products are required for mitogenesis and differentiation (By similarity).
Cellular Location	Nucleus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.