

TRIM9 Rabbit pAb

TRIM9 Rabbit pAb
Catalog # AP56574

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9C026
Reactivity	Mouse, Rat
Predicted	Human, Dog, Pig, Rabbit, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	79177
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TRIM9
Epitope Specificity	181-280/710
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasmic
SIMILARITY	Belongs to the TRIM/RBCC family. Contains 2 B box-type zinc fingers. Contains 1 B30.2/SPRY domain. Contains 1 COS domain. Contains 1 fibronectin type-III domain. Contains 1 RING-type zinc finger. Interacts with SNAP25.
SUBUNIT	
Post-translational modifications	Brain. Highly expressed in the cerebral cortex (at protein level). Severely decreased in the affected brain areas in Parkinson disease and dementia with Lewy bodies.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic bodies. Its function has not been identified. Alternate splicing of this gene generates two transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	114088
Other Names	E3 ubiquitin-protein ligase TRIM9, 2.3.2.27, RING finger protein 91, RING-type E3 ubiquitin transferase TRIM9, Tripartite motif-containing protein 9, TRIM9, KIAA0282, RNF91
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	TRIM9
Synonyms	KIAA0282, RNF91
Function	E3 ubiquitin-protein ligase which ubiquitinates itself in cooperation with an E2 enzyme UBE2D2/UBC4 and serves as a targeting signal for proteasomal degradation. May play a role in regulation of neuronal functions and may also participate in the formation or breakdown of abnormal inclusions in neurodegenerative disorders. May act as a regulator of synaptic vesicle exocytosis by controlling the availability of SNAP25 for the SNARE complex formation.
Cellular Location	Cytoplasm. Cell projection, dendrite. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle {ECO:0000250 UniProtKB:Q91ZY8}. Synapse {ECO:0000250 UniProtKB:Q91ZY8} Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:Q91ZY8}. Note=Enriched at synaptic terminals where it exists in a soluble form and a synaptic vesicle-associated form. Associated with the cytoskeleton (By similarity). Found in proximal dendrites of pyramidal neurons in the cerebral cortex and hippocampus, and Purkinje cells in the cerebellum (PubMed:20085810). {ECO:0000250 UniProtKB:Q91ZY8, ECO:0000269 PubMed:20085810}
Tissue Location	Brain. Highly expressed in the cerebral cortex (at protein level). Severely decreased in the affected brain areas in Parkinson disease and dementia with Lewy bodies

Background

The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic bodies. Its function has not been identified. Alternate splicing of this gene generates two transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

Images



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRIM9) Polyclonal Antibody, Unconjugated (AP56574) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.