

MCM8 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57013

Product Information

| | |
|--------------------------------|--|
| Application | IHC-P, IHC-F, IF, ICC |
| Primary Accession | Q9UJA3 |
| Reactivity | Dog, Bovine |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 93697 |
| Physical State | Liquid |
| Immunogen | KLH conjugated synthetic peptide derived from human MCM8 |
| Epitope Specificity | 301-400/840 |
| Isotype | IgG |
| Purity | affinity purified by Protein A |
| Buffer | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. |
| SUBCELLULAR LOCATION | Nucleus. Localizes to nuclear foci and colocalizes with RAD51. |
| SIMILARITY | Belongs to the MCM family. Contains 1 MCM domain. |
| Important Note | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |
| Background Descriptions | The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the mini-chromosome maintenance proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. This protein contains the central domain that is conserved among the mini-chromosome maintenance proteins. The encoded protein may interact with other mini-chromosome maintenance proteins and play a role in DNA replication. This gene may be associated with length of reproductive lifespan and menopause. Alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2013] |

Additional Information

| | |
|---------------------------|---|
| Gene ID | 84515 |
| Other Names | DNA helicase MCM8, 3.6.4.12, Minichromosome maintenance 8, MCM8, C20orf154 |
| Target/Specificity | Highest levels in placenta, lung and pancreas. Low levels in skeletal muscle and kidney. Expressed in various tumors with highest levels in colon and lung cancers. |
| Dilution | IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500 |

| | |
|----------------|---|
| Format | 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce |
| Storage | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |

Protein Information

| | |
|--------------------------|--|
| Name | MCM8 |
| Synonyms | C20orf154 |
| Function | Component of the MCM8-MCM9 complex, a complex involved in the repair of double-stranded DNA breaks (DBSs) and DNA interstrand cross- links (ICLs) by homologous recombination (HR) (PubMed: 23401855). Required for DNA resection by the MRE11-RAD50-NBN/NBS1 (MRN) complex by recruiting the MRN complex to the repair site and by promoting the complex nuclease activity (PubMed: 26215093). Probably by regulating the localization of the MNR complex, indirectly regulates the recruitment of downstream effector RAD51 to DNA damage sites including DBSs and ICLs (PubMed: 23401855). The MCM8-MCM9 complex is dispensable for DNA replication and S phase progression (PubMed: 23401855). However, may play a non-essential for DNA replication: may be involved in the activation of the prereplicative complex (pre-RC) during G(1) phase by recruiting CDC6 to the origin recognition complex (ORC) (PubMed: 15684404). Probably by regulating HR, plays a key role during gametogenesis (By similarity). Stabilizes MCM9 protein (PubMed: 23401855 , PubMed: 26215093). |
| Cellular Location | Nucleus. Chromosome. Note=Localizes to nuclear foci (PubMed: 26215093). Localizes to double-stranded DNA breaks (PubMed: 23401855). Binds chromatin throughout the cell cycle (PubMed: 15684404). |
| Tissue Location | Highest levels in placenta, lung and pancreas. Low levels in skeletal muscle and kidney. Expressed in various tumors with highest levels in colon and lung cancers |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.