

SLC10A1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5710B

Product Information

Application	WB, FC, E
Primary Accession	Q14973
Other Accession	NP_003040.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	316-343

Additional Information

Other Names	Sodium/bile acid cotransporter, Cell growth-inhibiting gene 29 protein, Na(+)/bile acid cotransporter, Na(+)/taurocholate transport protein, Sodium/taurocholate cotransporting polypeptide, Solute carrier family 10 member 1, SLC10A1, NTCP
Target/Specificity	This SLC10A1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 316-343 amino acids of human SLC10A1.
Dilution	WB~~1:2000 FC~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SLC10A1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Background

Sodium/bile acid cotransporters are integral membrane glycoproteins that participate in the enterohepatic circulation of bile acids. Two homologous transporters are involved in the reabsorption of bile acids, one absorbing from the intestinal lumen, the bile duct, and the kidney with an apical localization (SLC10A2; MIM 601295), and the other being found in the basolateral membranes of hepatocytes (SLC10A1).


References

Ho, R.H., et al. J. Biol. Chem. 279(8):7213-7222(2004)
Trauner, M., et al. Physiol. Rev. 83(2):633-671(2003)
Hallen, S., et al. Biochemistry 41(23):7253-7266(2002)
Shiao, T., et al. Genomics 69(2):203-213(2000)
Hagenbuch, B., et al. J. Clin. Invest. 93(3):1326-1331(1994)

Images



All lanes : Anti-SLC10A1 Antibody (C-term) at 1:1000 dilution
Lane 1: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 58 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



SLC10A1 Antibody (C-term) (Cat. #AP5710b) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Citations

- [Core-fucosylation plays a pivotal role in hepatitis B pseudo virus infection: a possible implication for HBV glycotherapy.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.