

Beta-phosphoglucomutase Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57140

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Reactivity	Lactococcus lactis
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from Lactococcus lactis Beta-phosphoglucomutase
Epitope Specificity	21-120/221
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasmic
SIMILARITY	Belongs to the HAD-like hydrolase superfamily. CbbY/CbbZ/Gph/YieH family.
Post-translational modifications	Autophosphorylated.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	<p>Beta-phosphoglucomutase is an enzyme that transfers a phosphoryl group on a glucose monomer from the 1' to the 6' position in the forward direction or the 6' to the 1' position in the reverse. Specifically, it converts Beta-D-glucose-1-phosphate to Beta-D-glucose-6-phosphate. This enzyme participates in both the breakdown and synthesis of glucose. Maltose metabolism in Lactococcus lactis involves the conversion of beta-glucose 1-phosphate to glucose 6-phosphate, a reaction which is reversibly catalysed by a maltose-inducible and glucose-repressible beta-phosphoglucomutase (beta-PGM). Alpha-PGM is expressed constitutively.</p> <p>Beta-phosphoglucomutase is a member of the haloacid dehalogenase superfamily of hydrolase enzymes. The enzyme from Lactococcus lactis has been extensively characterised including a remarkable crystal structure which traps the pentacoordinate transition state.</p>

Additional Information

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.