

NANS Rabbit pAb

NANS Rabbit pAb
Catalog # AP57352

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9NR45
Predicted	Human, Mouse, Rat, Chicken, Dog, Pig, Rabbit, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40308
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human NANS
Epitope Specificity	1-100/359
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 AFP-like domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes an enzyme that functions in the biosynthetic pathways of sialic acids. In vitro, the encoded protein uses N-acetylmannosamine 6-phosphate and mannose 6-phosphate as substrates to generate phosphorylated forms of N-acetylneuraminic acid (Neu5Ac) and 2-keto-3-deoxy-D-glycero-D-galacto-nononic acid (KDN), respectively; however, it exhibits much higher activity toward the Neu5Ac phosphate product. In insect cells, expression of this gene results in Neu5Ac and KDN production. This gene is related to the E. coli sialic acid synthase gene neuB, and it can partially restore sialic acid synthase activity in an E. coli neuB-negative mutant. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	54187
Other Names	N-acetylneuraminate-9-phosphate synthase, 2.5.1.57, 3-deoxy-D-glycero-D-galacto-nononate 9-phosphate synthase, 2.5.1.132, N-acetylneuraminic acid phosphate synthase, NANS, Sialic acid phosphate synthase, Sialic acid synthase, NANS (HGNC:19237), SAS
Target/Specificity	Ubiquitous.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	NANS (HGNC:19237)
Synonyms	SAS
Function	Catalyzes the condensation of phosphoenolpyruvate (PEP) and N-acetylmannosamine 6-phosphate (ManNAc-6-P) to synthesize N-acetylneuraminate-9-phosphate (Neu5Ac-9-P) (PubMed: 10749855). Also catalyzes the condensation of PEP and D-mannose 6-phosphate (Man-6-P) to produce 3-deoxy-D-glycero-beta-D-galacto-non-2-ulopyranosonate 9-phosphate (KDN-9-P) (PubMed: 10749855). Neu5Ac-9-P and KDN-9-P are the phosphorylated forms of sialic acids N-acetylneuraminic acid (Neu5Ac) and deaminoneuraminic acid (KDN), respectively (PubMed: 10749855). Required for brain and skeletal development (PubMed: 27213289).
Tissue Location	Ubiquitous..

Background

This gene encodes an enzyme that functions in the biosynthetic pathways of sialic acids. In vitro, the encoded protein uses N-acetylmannosamine 6-phosphate and mannose 6-phosphate as substrates to generate phosphorylated forms of N-acetylneuraminic acid (Neu5Ac) and 2-keto-3-deoxy-D-glycero-D-galacto-nononic acid (KDN), respectively; however, it exhibits much higher activity toward the Neu5Ac phosphate product. In insect cells, expression of this gene results in Neu5Ac and KDN production. This gene is related to the E. coli sialic acid synthase gene neuB, and it can partially restore sialic acid synthase activity in an E. coli neuB-negative mutant. [provided by RefSeq, Jul 2008]

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.