

PSG4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57527

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q00888
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47113
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human PSG4
Epitope Specificity	301-400/419
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
SIMILARITY	Belongs to the immunoglobulin superfamily. CEA family. Contains 3 Ig-like C2-type (immunoglobulin-like) domains. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The human pregnancy-specific glycoproteins (PSGs) are a family of proteins that are synthesized in large amounts by placental trophoblasts and released into the maternal circulation during pregnancy. Molecular cloning and analysis of several PSG genes has indicated that the PSGs form a subgroup of the carcinoembryonic antigen (CEA) gene family, which belongs to the immunoglobulin superfamily of genes. Members of the CEA family consist of a single N domain, with structural similarity to the immunoglobulin variable domains, followed by a variable number of immunoglobulin constant-like A and/or B domains. Most PSGs have an arg-gly-asp (RGD) motif, which has been shown to function as an adhesion recognition signal for several integrins, in the N-terminal domain (summary by Teglund et al., 1994 [PubMed 7851896]). For additional general information about the PSG gene family, see PSG1 (MIM 176390).[supplied by OMIM, Oct 2009]

Additional Information


Gene ID	5672
Other Names	Pregnancy-specific beta-1-glycoprotein 4, PS-beta-G-4, PSBG-4, Pregnancy-specific glycoprotein 4, Pregnancy-specific beta-1-glycoprotein 9, PS-beta-G-9, PSBG-9, Pregnancy-specific glycoprotein 9, PSG4, CGM4, PSG9
Target/Specificity	PSBG are produced in high quantity during pregnancy.

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	PSG4
Synonyms	CGM4, PSG9
Cellular Location	Secreted.

Images




Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PSG4) Polyclonal Antibody, Unconjugated (AP57527) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample:

MDA-MB-231(Human) Cell Lysate at 30 ug
 293T(Human) Cell Lysate at 30 ug
 Primary: Anti-PSG4 (AP57527) at 1/300 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 47 kD
 Observed band size: 42/57 kD



Sample:

293T(Human) Lysate at 30 ug
 Primary: Anti- PSG4 (AP57527) at 1/300 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 47 kD
 Observed band size: 57/67 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.