

eIF3A Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58097

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q14152
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	166569
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human eIF3A
Epitope Specificity	801-900/1382
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm.
SIMILARITY	Belongs to the eIF-3 subunit A family.Contains 1 PCI domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA ⁱ and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation.

Additional Information

Gene ID	8661
Other Names	Eukaryotic translation initiation factor 3 subunit A {ECO:0000255 HAMAP-Rule:MF_03000}, eIF3a {ECO:0000255 HAMAP-Rule:MF_03000}, Eukaryotic translation initiation factor 3 subunit 10 {ECO:0000255 HAMAP-Rule:MF_03000}, eIF-3-theta {ECO:0000255 HAMAP-Rule:MF_03000}, eIF3 p167, eIF3 p180, eIF3 p185, EIF3A {ECO:0000255 HAMAP-Rule:MF_03000}
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name

EIF3A {ECO:0000255 | HAMAP-Rule:MF_03000}


Function

RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:[17581632](#), PubMed:[25849773](#)). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl- tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:[11169732](#), PubMed:[17581632](#)). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem- loop binding to exert either translational activation or repression (PubMed:[25849773](#), PubMed:[27462815](#)).


Cellular Location

Cytoplasm {ECO:0000255 | HAMAP-Rule:MF_03000, ECO:0000269 | PubMed:9150439}

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (eIF3A) Polyclonal Antibody, Unconjugated (AP58097) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Sample:

Testis (Mouse) Lysate at 40 ug

HeLa(Human) Cell Lysate at 30 ug

Primary: Anti- eIF3A (AP58097) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 166 kD

Observed band size: 166 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.