

P2Y2 Rabbit pAb

P2Y2 Rabbit pAb Catalog # AP58129

Product Information

Application WB, IHC-P, IHC-F, IF, E

Primary Accession P41231
Reactivity Human

Predicted Mouse, Rat, Dog, Horse, Rabbit

Host Rabbit
Clonality Polyclonal
Calculated MW 42273
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human P2Y2R

Epitope Specificity 21-120/377

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cell membrane; Multi-pass membrane protein.

SIMILARITY Belongs to the G-protein coupled receptor 1 family.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions The product of this gene belongs to the family of G-protein coupled

receptors. This family has several receptor subtypes with different pharmacological selectivity, which overlaps in some cases, for various adenosine and uridine nucleotides. This receptor is responsive to both adenosine and uridine nucleotides. It may participate in control of the cell cycle of endometrial carcinoma cells. Three transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq].

Additional Information

Gene ID 5029

Other Names P2Y purinoceptor 2, P2Y2, ATP receptor, P2U purinoceptor 1, P2U1, P2U

receptor 1, Purinergic receptor, P2RY2, P2RU1

Target/Specificity Spleen, testis, kidney, liver, lung, heart and brain.

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000

-10000

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name P2RY2

Synonyms P2RU1

Function Receptor for ATP and UTP coupled to G-proteins that activate a

phosphatidylinositol-calcium second messenger system. The affinity range is

UTP = ATP > ATP-gamma-S >> 2-methylthio-ATP = ADP.

Cellular Location Cell membrane; Multi-pass membrane protein.

Tissue Location Spleen, testis, kidney, liver, lung, heart and brain

Background

The product of this gene belongs to the family of G-protein coupled receptors. This family has several receptor subtypes with different pharmacological selectivity, which overlaps in some cases, for various adenosine and uridine nucleotides. This receptor is responsive to both adenosine and uridine nucleotides. It may participate in control of the cell cycle of endometrial carcinoma cells. Three transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq].

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.