

SDF2L1 Rabbit pAb

SDF2L1 Rabbit pAb
Catalog # AP58210

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9HCN8
Reactivity	Mouse
Predicted	Human, Dog, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23598
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SDF2L1
Epitope Specificity	121-221/221
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4.
SUBCELLULAR LOCATION	Endoplasmic reticulum lumen.
SIMILARITY	Contains 3 MIR domains.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	SDF2L1 (Stromal Cell Derived Factor 2 Like 1) is a Protein Coding gene. GO annotations related to this gene include chaperone binding and misfolded protein binding. An important paralog of this gene is SDF2.

Additional Information

Gene ID	23753
Other Names	Stromal cell-derived factor 2-like protein 1, SDF2-like protein 1, PWP1-interacting protein 8, SDF2L1
Target/Specificity	Ubiquitously expressed with high expression in testis, moderate expression in the pancreas, spleen, prostate, small intestine and colon. Very low expression is seen in brain and skeletal muscle.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SDF2L1
Cellular Location	Endoplasmic reticulum lumen {ECO:0000255 PROSITE- ProRule:PRU10138}
Tissue Location	Ubiquitously expressed with high expression in testis, moderate expression in the pancreas, spleen, prostate, small intestine and colon. Very low expression is seen in brain and skeletal muscle.

Background

SDF2L1 (Stromal Cell Derived Factor 2 Like 1) is a Protein Coding gene. GO annotations related to this gene include chaperone binding and misfolded protein binding. An important paralog of this gene is SDF2.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.