

# SLC9A3R1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP5853c

#### **Product Information**

**Application** WB, FC, E **Primary Accession** 014745

**Reactivity** Human, Rat, Mouse

**Predicted** Bovine, Monkey, Mouse, Rat

HostRabbitClonalityPolyclonalIsotypeRabbit IgGClone NamesRB21816Calculated MW38868Antigen Region168-197

## **Additional Information**

**Gene ID** 9368

Other Names Na(+)/H(+) exchange regulatory cofactor NHE-RF1, NHERF-1,

Ezrin-radixin-moesin-binding phosphoprotein 50, EBP50, Regulatory cofactor of Na(+)/H(+) exchanger, Sodium-hydrogen exchanger regulatory factor 1, Solute carrier family 9 isoform A3 regulatory factor 1, SLC9A3R1, NHERF,

NHERF1

**Target/Specificity** This SLC9A3R1 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 168-197 amino acids of human

SLC9A3R1.

**Dilution** WB~~1:2000 FC~~1:10~50 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** SLC9A3R1 Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

### **Protein Information**

Name NHERF1 ( HGNC:11075)

#### **Synonyms**

NHERF, SLC9A3R1

#### **Function**

Scaffold protein that connects plasma membrane proteins with members of the ezrin/moesin/radixin family and thereby helps to link them to the actin cytoskeleton and to regulate their surface expression. Necessary for recycling of internalized ADRB2. Was first known to play a role in the regulation of the activity and subcellular location of SLC9A3. Necessary for cAMP-mediated phosphorylation and inhibition of SLC9A3. May enhance Wnt signaling. May participate in HTR4 targeting to microvilli (By similarity). Involved in the regulation of phosphate reabsorption in the renal proximal tubules. Involved in sperm capacitation. May participate in the regulation of the chloride and bicarbonate homeostasis in spermatozoa.


#### **Cellular Location**

Cytoplasm. Apical cell membrane. Endomembrane system; Peripheral membrane protein. Cell projection, filopodium. Cell projection, ruffle. Cell projection, microvillus. Note=Translocates from the cytoplasm to the apical cell membrane in a PODXL-dependent manner. Colocalizes with CFTR at the midpiece of sperm tail (By similarity). Colocalizes with actin in microvilli-rich apical regions of the syncytiotrophoblast. Found in microvilli, ruffling membrane and filopodia of HeLa cells. Present in lipid rafts of T-cells.


#### **Tissue Location**

Detected in liver, kidney, pancreas, prostate, spleen, small intestine and placenta, in particular in the syncytiotrophoblast.

## **Images**




All lanes: Anti-SLC9A3R1 Antibody (Center) at 1:2000 dilution Lane 1: HepG2 whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: T47D whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 39 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



SLC9A3R1 Antibody (Center) (Cat. #AP5853c) western blot analysis in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the SLC9A3R1 antibody detected the SLC9A3R1 protein (arrow).

SLC9A3R1 Antibody (Center) (Cat. #AP5853c) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



## **Citations**

• Cell cycle dependent association of EBP50 with protein phosphatase 2A in endothelial cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.