

# DMTF1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58544

## Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">Q9Y222</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	84471

## Additional Information

Gene ID	9988
Other Names	Cyclin-D-binding Myb-like transcription factor 1, hDMTF1, Cyclin-D-interacting Myb-like protein 1, hDMP1, DMTF1, DMP1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

Name	DMTF1
Synonyms	DMP1
Function	Transcriptional activator which activates the CDKN2A/ARF locus in response to Ras-Raf signaling, thereby promoting p53/TP53- dependent growth arrest (By similarity). Binds to the consensus sequence 5'-CCCG[GT]ATGT-3' (By similarity). Isoform 1 may cooperate with MYB to activate transcription of the ANPEP gene. Isoform 2 may antagonize transcriptional activation by isoform 1.
Cellular Location	Nucleus {ECO:0000255   PROSITE-ProRule:PRU00625, ECO:0000269   PubMed:17936562}
Tissue Location	Expressed at relatively low levels in colonic mucosa, ovary, peripheral leukocytes, prostate and small intestine, and at higher levels in spleen, testis and thymus. Expressed in multiple regions of the brain and CNS including amygdala, caudate, corpus callosum, hippocampus, substantia nigra and

subthalamic nucleus Isoform 1 is the predominant isoform in monocytes, macrophages and neutrophils, isoform 2 is most strongly expressed in peripheral blood leukocytes and quiescent CD34 positive cells, and isoform 3 is expressed at low levels in all hematopoietic cell types. Expression is frequently reduced in non-small-cell lung carcinomas (NSCLC) due to hemizygous gene deletion, strongly suggesting that this locus is haploinsufficient for tumor suppression. Loss of this locus frequently occurs in tumors which retain wild-type CDKN2A/ARF and p53/TP53 loci Hemizygous gene deletion has also been observed in leukemic blasts from patients with abnormalities of the long arm of chromosome 7

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.