

PPARG Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5863a

Product Information

Application	WB, FC, E
Primary Accession	P37231
Other Accession	NP_619725.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB22139
Calculated MW	57620
Antigen Region	1-30

Additional Information

Gene ID	5468
Other Names	Peroxisome proliferator-activated receptor gamma, PPAR-gamma, Nuclear receptor subfamily 1 group C member 3, PPARG, NR1C3
Target/Specificity	This PPARG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human PPARG.
Dilution	WB~~1:2000 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PPARG Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PPARG
Synonyms	NR1C3
Function	Ligand-activated transcription factor that forms obligate heterodimers with

the retinoic acid receptor and acts as a key regulator of biological processes, such as adipocyte differentiation, lipid metabolism, glucose homeostasis and beta-oxidation of fatty acids (PubMed:[16150867](#), PubMed:[20829347](#), PubMed:[23525231](#), PubMed:[8702406](#), PubMed:[8706692](#), PubMed:[9065481](#)). Activated by lipid ligands: binds peroxisome proliferators, such as hypolipidemic drugs, and fatty acids, such as prostaglandin J2 metabolites (PubMed:[16150867](#), PubMed:[20829347](#), PubMed:[23525231](#), PubMed:[8702406](#), PubMed:[8706692](#), PubMed:[9065481](#)). Ligand-binding results in a conformational change in the receptor, promoting dissociation of repressors and recruitment of coactivators, and subsequent activation of target gene expression (PubMed:[16150867](#), PubMed:[20829347](#), PubMed:[23525231](#), PubMed:[8702406](#), PubMed:[8706692](#), PubMed:[9065481](#)). Specifically binds to DNA specific PPAR response elements (PPRE) and modulates the transcription of its target genes, such as acyl-CoA oxidase (By similarity). Acts as a critical regulator of gut homeostasis by suppressing NF-kappa-B-mediated pro-inflammatory responses (PubMed:[20829347](#)). Plays a role in the regulation of cardiovascular circadian rhythms by regulating the transcription of BMAL1 in the blood vessels (By similarity).


Cellular Location

Nucleus. Cytoplasm Note=Redistributed from the nucleus to the cytosol through a MAP2K1/MEK1-dependent manner (PubMed:[17101779](#)). NOCT enhances its nuclear translocation (By similarity). {ECO:[0000250](#) | UniProtKB:P37238, ECO:[0000269](#) | PubMed:[17101779](#)}


Tissue Location

Highest expression in adipose tissue. Lower in skeletal muscle, spleen, heart and liver. Also detectable in placenta, lung and ovary.

Images



Overlay histogram showing HeLa cells stained with AP5863a(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP5863a, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1μg/1x10^6 cells) used under the same conditions. Acquisition of >10, 000 events was performed.



All lanes : Anti-PPARG Antibody (N-term) at 1:2000 dilution Lane 1: Human fetal liver lysate Lane 2: Human lung lysate Lane 3: Human placenta lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 58 kDa Blocking/Dilution buffer: 5% NFDM/TBST.