

beta Defensin 3 Rabbit pAb

beta Defensin 3 Rabbit pAb Catalog # AP58656

Product Information

Primary Accession
Reactivity
Host
Clonality
Calculated MW
Physical State
R81534
Human
Puman
Rabbit
Polyclonal
7697
Liquid

Immunogen KLH conjugated synthetic peptide derived from human beta Defensin 3

Epitope Specificity 23-67/67 **Isotype** IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Secreted.

SIMILARITY Belongs to the beta-defensin family.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions Exhibits antimicrobial activity against Gram-positive bacteria S.aureus and

S.pyogenes, Gram-negative bacteria P.aeruginosa and E.coli and the yeast C.albicans. Kills multiresistant S.aureus and vancomycin-resistent E.faecium.

No significant hemolytic activity was observed.

Additional Information

Gene ID 414325;55894

Other Names Beta-defensin 103, Beta-defensin 3, BD-3, DEFB-3, HBD3, hBD-3, Defensin,

beta 103, Defensin-like protein, DEFB103A, BD3, DEFB103, DEFB3

Target/Specificity Highly expressed in skin and tonsils, and to a lesser extent in trachea, uterus,

kidney, thymus, adenoid, pharynx and tongue. Low expression in salivary gland, bone marrow, colon, stomach, polyp and larynx. No expression in

small intestine.

Dilution Flow-Cyt=1 □g/Test

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name DEFB103A

Synonyms BD3, DEFB103, DEFB3

Function Exhibits antimicrobial activity against Gram-positive bacteria S.aureus and

S.pyogenes, Gram-negative bacteria P.aeruginosa and E.coli and the yeast C.albicans. Kills multiresistant S.aureus and vancomycin-resistant E.faecium.

No significant hemolytic activity was observed.

Cellular Location Secreted.

Tissue Location Highly expressed in skin and tonsils, and to a lesser extent in trachea, uterus,


kidney, thymus, adenoid, pharynx and tongue. Low expression in salivary gland, bone marrow, colon, stomach, polyp and larynx. No expression in

small intestine


Background

Exhibits antimicrobial activity against Gram-positive bacteria S.aureus and S.pyogenes, Gram-negative bacteria P.aeruginosa and E.coli and the yeast C.albicans. Kills multiresistant S.aureus and vancomycin-resistent E.faecium. No significant hemolytic activity was observed.

Images



Blank control(blue): Hep G2 Cells(fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice). Primary Antibody: Rabbit Anti-beta Defensin 3/FITC Conjugated antibody (AP58656-FITC), Dilution: 1 µg in 100 µL 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG/FITC(orange) ,used under the same conditions.



Blank control(blue): Hep G2 Cells(fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice).

Primary Antibody: Rabbit Anti-beta Defensin 3/FITC
Conjugated antibody (AP58656-FITC), Dilution: 1 µg in 100 µL 1X PBS containing 0.5% BSA;
Isotype Control Antibody: Rabbit IgG/FITC(orange), used under the same conditions.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.