

# **AMSH Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58738

#### **Product Information**

**Application** IHC-P, IHC-F, IF, E

Primary Accession

Reactivity

Rost

Clonality

Calculated MW

Physical State

O95630

Rat, Dog

Rabbit

Polyclonal

48077

Liquid

Immunogen KLH conjugated synthetic peptide derived from human AMSH

Epitope Specificity 331-420/424

**Isotype** IgG

**Purity** affinity purified by Protein A

**Buffer** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus. Membrane; Peripheral membrane protein. Cytoplasm. Early

endosome.

**SIMILARITY** Belongs to the peptidase M67C family.Contains 1 MPN (JAB/Mov34) domain. **Post-translational** Phosphorylated after BMP type I receptor activation.Ubiquitinated by SMURF2

**modifications** in the presence of RNF11.

**Important Note** This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

**Background Descriptions** Zinc metalloprotease that specifically cleaves 'Lys-63'-linked polyubiquitin

chains. Does not cleave 'Lys-48'-linked polyubiquitin chains (By similarity). Functions at the endosome and is able to oppose the ubiquitin-dependent sorting of receptors to lysosomes. Plays a role in signal transduction for cell growth and MYC induction mediated by IL-2 and GM-CSF. Potentiates BMP (bone morphogenetic protein) signaling by antagonizing the inhibitory action

of SMAD6 and SMAD7.

#### **Additional Information**

**Gene ID** 10617

**Other Names** STAM-binding protein, 3.4.19.-, Associated molecule with the SH3 domain of

STAM, Endosome-associated ubiquitin isopeptidase, STAMBP, AMSH

**Target/Specificity** Ubiquitously expressed.

**Dilution** IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### **Protein Information**

Name STAMBP

**Synonyms** AMSH {ECO:0000303 | PubMed:10383417}

**Function** Zinc metalloprotease that specifically cleaves 'Lys-63'- linked polyubiquitin

chains (PubMed: 15314065, PubMed: 23542699, PubMed: 34425109). Does not cleave 'Lys-48'-linked polyubiquitin chains (PubMed: 15314065). Plays a role in signal transduction for cell growth and MYC induction mediated by IL-2 and GM-CSF (PubMed: 10383417). Potentiates BMP (bone morphogenetic protein) signaling by antagonizing the inhibitory action of SMAD6 and SMAD7

(PubMed:<u>11483516</u>). Has a key role in regulation of cell surface

receptor-mediated endocytosis and ubiquitin-dependent sorting of receptors to lysosomes (PubMed:15314065, PubMed:17261583). Endosomal localization

of STAMBP is required for efficient EGFR degradation but not for its internalization (PubMed: 15314065, PubMed: 17261583). Involved in the negative regulation of PI3K-AKT-mTOR and RAS-MAP signaling pathways


(PubMed:23542699).

**Cellular Location** Nucleus. Membrane; Peripheral membrane protein. Cytoplasm. Early

endosome

**Tissue Location** Ubiquitously expressed.

## **Images**




Tissue/cell: mouse embryo tissue; 4%
Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3%
Hydrogen peroxide for 30min; Blocking buffer (normal

Unconjugated(AP58738) 1:200, overnight at 4°C, followed

Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-AMSH/STAMBP Polyclonal Antibody,

by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (AMSH) Polyclonal Antibody, Unconjugated (AP58738) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.