

RNF98 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP59163

Product Information

| | |
|--------------------------------|---|
| Application | IHC-P, IHC-F, IF, E |
| Primary Accession | Q9NQ86 |
| Reactivity | Rat, Pig, Bovine |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 83013 |
| Physical State | Liquid |
| Immunogen | KLH conjugated synthetic peptide derived from human TRIM36/RNF98 |
| Epitope Specificity | 201-300/728 |
| Isotype | IgG |
| Purity | affinity purified by Protein A |
| Buffer | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. |
| SUBCELLULAR LOCATION | Cytoplasm (By similarity). Cytoplasmic vesicle, secretory vesicle, acrosome (By similarity). Cytoplasm, cytoskeleton (By similarity). Note=Found in the acrosomal region of elongated spermatids and mature sperm (By similarity). |
| SIMILARITY | Belongs to the TRIM/RBCC family. Contains 2 B box-type zinc fingers. Contains 1 B30.2/SPRY domain. Contains 1 COS domain. Contains 1 fibronectin type-III domain. Contains 1 RING-type zinc finger. |
| SUBUNIT | Interacts with CENPH (By similarity). |
| Important Note | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |
| Background Descriptions | TRIM36 (tripartite motif-containing 36), also known as RNF98 (RING finger protein 98), HAPRIN (haploid germ cell-specific RBCC protein) or RBCC728, is a 728 amino acid protein that belongs to the TRIM/RBCC (Ring finger, B box, coiled-coil) family. Predominantly expressed in prostate, testis and brain with weak expression in heart, kidney and lung, TRIM36 contains two B box-type zinc fingers, a SPRY domain, a coiled-coil domain, a fibronectin type-III domain and a RING-type zinc finger; a motif that has zinc-chelating activity and is involved in mediating protein-protein and protein-DNA interactions. Localizing to the cytoplasm and the acrosomal region of germ cells and mature sperm, TRIM36 is believed to play a role in the acrosome reaction and fertilization. In addition, TRIM36 is overexpressed in prostate cancer, suggesting a possible role for TRIM36 in prostate tumorigenesis. |

Additional Information

| | |
|--------------------|--|
| Gene ID | 55521 |
| Other Names | E3 ubiquitin-protein ligase TRIM36, 2.3.2.27, Tripartite motif-containing protein 36, Zinc-binding protein Rbcc728, TRIM36, RBCC728, RNF98 |

| | |
|---------------------------|---|
| Target/Specificity | Highly expressed in testis, prostate and brain. Weakly expressed in kidney, lung and heart. |
| Dilution | IHC-P=1:100-500,IHC-F=1:100-500,IF=1:50-200,ELISA=1:5000-10000 |
| Format | 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce |
| Storage | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |

Protein Information

| | |
|--------------------------|--|
| Name | TRIM36 |
| Synonyms | RBCC728, RNF98 |
| Function | E3 ubiquitin-protein ligase which mediates ubiquitination and subsequent proteasomal degradation of target proteins. Involved in chromosome segregation and cell cycle regulation (PubMed: 28087737). May play a role in the acrosome reaction and fertilization. |
| Cellular Location | Cytoplasm. Cytoplasmic vesicle, secretory vesicle, acrosome {ECO:0000250 UniProtKB:Q80WG7}. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:Q80WG7}. Note=Found in the acrosomal region of elongated spermatids and mature sperm. {ECO:0000250 UniProtKB:Q80WG7} |
| Tissue Location | Highly expressed in testis, prostate and brain (PubMed:15145053). Weakly expressed in kidney, lung and heart (PubMed:15145053). Expressed in fetal tissues (PubMed:28087737) |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.