

# Anti-Arginase 2 Antibody

Rabbit polyclonal antibody to Arginase 2  
Catalog # AP59482

## Product Information

---

<b>Application</b>	WB, IP
<b>Primary Accession</b>	<a href="#">P78540</a>
<b>Reactivity</b>	Human, Monkey
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	38578

## Additional Information

---

<b>Gene ID</b>	384
<b>Other Names</b>	Arginase-2 mitochondrial; Kidney-type arginase; Non-hepatic arginase; Type II arginase
<b>Target/Specificity</b>	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Arginase 2. The exact sequence is proprietary.
<b>Dilution</b>	WB~~WB (1/500 - 1/1000), IP (1/10 - 1/100) IP~~N/A
<b>Format</b>	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
<b>Storage</b>	Store at -20 °C. Stable for 12 months from date of receipt

## Protein Information

---

<b>Name</b>	ARG2
<b>Function</b>	May play a role in the regulation of extra-urea cycle arginine metabolism and also in down-regulation of nitric oxide synthesis. Extrahepatic arginase functions to regulate L-arginine bioavailability to nitric oxid synthase (NOS). Arginine metabolism is a critical regulator of innate and adaptive immune responses. Seems to be involved in negative regulation of the survival capacity of activated CD4(+) and CD8(+) T cells (PubMed: <a href="#">27745970</a> ). May suppress inflammation- related signaling in asthmatic airway epithelium (PubMed: <a href="#">27214549</a> ). May contribute to the immune evasion of H.pylori by restricting M1 macrophage activation and polyamine metabolism (By similarity). In fetal dendritic cells may play a role in promoting immune suppression and T cell TNF production during gestation (PubMed: <a href="#">28614294</a> ). Regulates RPS6KB1 signaling, which promotes endothelial cell senescence and inflammation and implicates NOS3/eNOS dysfunction (PubMed: <a href="#">22928666</a> ). Can inhibit endothelial autophagy independently of its enzymatic activity

implicating mTORC2 signaling (PubMed:[25484082](#)). Involved in vascular smooth muscle cell senescence and apoptosis independently of its enzymatic activity (PubMed:[23832324](#)). Since NOS is found in the penile corpus cavernosum smooth muscle, the clitoral corpus cavernosum and the vagina, arginase-2 plays a role in both male and female sexual arousal (PubMed:[12859189](#)).

**Cellular Location**

Mitochondrion.

**Tissue Location**

Expressed most strongly in kidney and prostate, much less strongly in the brain, skeletal muscle, placenta, lung, mammary gland, macrophage, uterus, testis and gut, but apparently not in the liver, heart and pancreas. Expressed in activated T cells (PubMed:[27745970](#)).


**Background**

---

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Arginase 2. The exact sequence is proprietary.

**Images**

---



Western blot analysis of Arginase 2 expression in HEK293T (A), HeLa (B), DLD (C) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.