

Anti-SMAD1 (pS187) Antibody

Rabbit polyclonal antibody to SMAD1 (pS187)

Catalog # AP59613

Product Information

| | |
|--------------------------|----------------------------------|
| Application | WB, IP, IHC |
| Primary Accession | Q15797 |
| Other Accession | P70340 |
| Reactivity | Human, Mouse, Rat, Chicken, SARS |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 52260 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 4086 |
| Other Names | BSP1; MADH1; MADR1; Mothers against decapentaplegic homolog 1; MAD homolog 1; Mothers against DPP homolog 1; JV4-1; Mad-related protein 1; SMAD family member 1; SMAD 1; Smad1; hSMAD1; Transforming growth factor-beta-signaling protein 1; BSP-1 |
| Target/Specificity | KLH-conjugated synthetic peptide encompassing a sequence within the center region of human SMAD1. The exact sequence is proprietary. |
| Dilution | WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IP (1/10 - 1/100) IP~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IP (1/10 - 1/100) |
| Format | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide. |
| Storage | Store at -20 °C. Stable for 12 months from date of receipt |

Protein Information

| | |
|-----------------|--|
| Name | SMAD1 (HGNC:6767) |
| Synonyms | BSP1, MADH1, MADR1 |
| Function | Transcriptional modulator that plays a role in various cellular processes, including embryonic development, cell differentiation, and tissue homeostasis (PubMed: 9335504). Upon BMP ligand binding to their receptors at the cell surface, is phosphorylated by activated type I BMP receptors (BMPRIs) and associates with SMAD4 to form a heteromeric complex which translocates into the nucleus acting as transcription factor (PubMed: 33667543). In turn, the hetero-trimeric complex recognizes cis-regulatory elements containing Smad Binding Elements (SBEs) to modulate the outcome of the signaling |

network (PubMed:[33667543](#)). SMAD1/OAZ1/PSMB4 complex mediates the degradation of the CREBBP/EP300 repressor SNIP1. Positively regulates BMP4-induced expression of odontogenic development regulator MSX1 following IPO7-mediated nuclear import (By similarity).

Cellular Location

Cytoplasm. Nucleus Note=Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4 (PubMed:15647271). Co-localizes with LEMD3 at the nucleus inner membrane (PubMed:15647271). Exported from the nucleus to the cytoplasm when dephosphorylated (By similarity) {ECO:0000250 | UniProtKB:P70340, ECO:0000269 | PubMed:15647271}


Tissue Location

Ubiquitous. Highest expression seen in the heart and skeletal muscle


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human SMAD1. The exact sequence is proprietary.

Images



Western blot analysis of SMAD1 (pS187) expression in A375 (A) whole cell lysates.



Immunohistochemical analysis of SMAD1 (pS187) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.