

Anti-PPAR gamma (pS112) Antibody

Rabbit polyclonal antibody to PPAR gamma (pS112)

Catalog # AP59666

Product Information

Application	WB
Primary Accession	P37231
Other Accession	P37238
Reactivity	Human, Mouse, Rat, Rabbit, Drosophila
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57620

Additional Information

Gene ID	5468
Other Names	NR1C3; Peroxisome proliferator-activated receptor gamma; PPAR-gamma; Nuclear receptor subfamily 1 group C member 3
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human PPAR gamma. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C. Stable for 12 months from date of receipt

Protein Information

Name	PPARG
Synonyms	NR1C3
Function	Ligand-activated transcription factor that forms obligate heterodimers with the retinoic acid receptor and acts as a key regulator of biological processes, such as adipocyte differentiation, lipid metabolism, glucose homeostasis and beta-oxidation of fatty acids (PubMed: 16150867 , PubMed: 20829347 , PubMed: 23525231 , PubMed: 8702406 , PubMed: 8706692 , PubMed: 9065481). Activated by lipid ligands: binds peroxisome proliferators, such as hypolipidemic drugs, and fatty acids, such as prostaglandin J2 metabolites (PubMed: 16150867 , PubMed: 20829347 , PubMed: 23525231 , PubMed: 8702406 , PubMed: 8706692 , PubMed: 9065481). Ligand-binding results in a conformational change in the receptor, promoting dissociation of repressors and recruitment of coactivators, and subsequent activation of

target gene expression (PubMed:[16150867](#), PubMed:[20829347](#), PubMed:[23525231](#), PubMed:[8702406](#), PubMed:[8706692](#), PubMed:[9065481](#)). Specifically binds to DNA specific PPAR response elements (PPRE) and modulates the transcription of its target genes, such as acyl-CoA oxidase (By similarity). Acts as a critical regulator of gut homeostasis by suppressing NF-kappa-B-mediated pro-inflammatory responses (PubMed:[20829347](#)). Plays a role in the regulation of cardiovascular circadian rhythms by regulating the transcription of BMAL1 in the blood vessels (By similarity).

Cellular Location

Nucleus. Cytoplasm Note=Redistributed from the nucleus to the cytosol through a MAP2K1/MEK1-dependent manner (PubMed:17101779). NOCT enhances its nuclear translocation (By similarity). {ECO:0000250 | UniProtKB:P37238, ECO:0000269 | PubMed:17101779}


Tissue Location

Highest expression in adipose tissue. Lower in skeletal muscle, spleen, heart and liver. Also detectable in placenta, lung and ovary.

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human PPAR gamma. The exact sequence is proprietary.

Images



Western blot analysis of PPAR gamma (pS112) expression in mouse heart (A), rat heart (B), H9C2 (C), AML12 (D), A549 (E), A2780 (F), LO2 (G) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.