

Anti-STK17B Antibody

Rabbit polyclonal antibody to STK17B Catalog # AP59775

Product Information

Application WB, IHC
Primary Accession 094768
Other Accession Q88G48

Reactivity Human, Mouse, Rat, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 42344

Additional Information

Gene ID 9262

Other Names DRAK2; Serine/threonine-protein kinase 17B; DAP kinase-related

apoptosis-inducing protein kinase 2

Target/Specificity Recognizes endogenous levels of STK17B protein.

Dilution WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200) IHC~~WB (1/500 - 1/1000), IHC

(1/100 - 1/200)

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name STK17B

Synonyms DRAK2

Function Phosphorylates myosin light chains (By similarity). Acts as a positive

regulator of apoptosis.

Cellular Location Nucleus. Cell membrane. Endoplasmic reticulum-Golgi intermediate

compartment. Note=Colocalizes with STK17B at the plasma membrane.


Tissue Location Highly expressed in placenta, lung, pancreas. Lower levels in heart, brain,

liver, skeletal muscle and kidney


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human STK17B. The exact sequence is proprietary.

Images



Western blot analysis of STK17B expression in MCF7 (A), K562 (B), HEK293T (C), mouse brain (D), rat brain (E) whole cell lysates.



Immunohistochemical analysis of STK17B staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.