

Anti-TXNRD2 Antibody

Rabbit polyclonal antibody to TXNRD2
Catalog # AP59800

Product Information

Application	WB, IHC
Primary Accession	Q9NNW7
Reactivity	Human, Mouse, Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56507

Additional Information

Gene ID	10587
Other Names	KIAA1652; TRXR2; Thioredoxin reductase 2 mitochondrial; Selenoprotein Z; SelZ; TR-beta; Thioredoxin reductase TR3
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human TXNRD2. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C. Stable for 12 months from date of receipt

Protein Information


Name	TXNRD2 (HGNC:18155)
Synonyms	KIAA1652, TRXR2
Function	Involved in the control of reactive oxygen species levels and the regulation of mitochondrial redox homeostasis (PubMed: 24601690). Maintains thioredoxin in a reduced state. May play a role in redox- regulated cell signaling.
Cellular Location	Mitochondrion.
Tissue Location	Highly expressed in the prostate, ovary, liver, testis, uterus, colon and small intestine. Intermediate levels in brain, skeletal muscle, heart and spleen. Low levels in placenta, pancreas, thymus and peripheral blood leukocytes. According to PubMed:10608886, high levels in kidney, whereas according to

PubMed:9923614, levels are low. High expression is observed in the adrenal cortex (PubMed:24601690).


Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human TXNRD2. The exact sequence is proprietary.

Images



Western blot analysis of TXNRD2 expression in HEK293T (A), HGC27 (B), mouse lung (C), mouse liver (D), rat lung (E), rat liver (F) whole cell lysates.



Immunohistochemical analysis of TXNRD2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.