

Anti-LATS1 Antibody

Rabbit polyclonal antibody to LATS1

Catalog # AP59945

Product Information

Application	WB
Primary Accession	Q95835
Other Accession	Q8BYR2
Reactivity	Human, Mouse, Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	126870

Additional Information

Gene ID	9113
Other Names	WARTS; Serine/threonine-protein kinase LATS1; Large tumor suppressor homolog 1; WARTS protein kinase; h-warts
Target/Specificity	Recognizes endogenous levels of LATS1 protein.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	LATS1 {ECO:0000312 EMBL:AAD16882.1}
Function	Negative regulator of YAP1 in the Hippo signaling pathway that plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis (PubMed: 10518011 , PubMed: 10831611 , PubMed: 18158288 , PubMed: 26437443 , PubMed: 28068668). The core of this pathway is composed of a kinase cascade wherein STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ (PubMed: 18158288 , PubMed: 26437443 , PubMed: 28068668). Phosphorylation of YAP1 by LATS1 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration (PubMed: 18158288 , PubMed: 26437443 , PubMed: 28068668). Acts as a tumor suppressor which plays a critical role in maintenance of ploidy through its actions in both

mitotic progression and the G1 tetraploidy checkpoint (PubMed:[15122335](#), PubMed:[19927127](#)). Negatively regulates G2/M transition by down-regulating CDK1 kinase activity (PubMed:[9988268](#)). Involved in the control of p53 expression (PubMed:[15122335](#)). Affects cytokinesis by regulating actin polymerization through negative modulation of LIMK1 (PubMed:[15220930](#)). May also play a role in endocrine function. Plays a role in mammary gland epithelial cell differentiation, both through the Hippo signaling pathway and the intracellular estrogen receptor signaling pathway by promoting the degradation of ESR1 (PubMed:[28068668](#)). Acts as an activator of the NLRP3 inflammasome by mediating phosphorylation of 'Ser-265' of NLRP3 following NLRP3 palmitoylation, promoting NLRP3 activation by NEK7 (PubMed:[39173637](#)).

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Midbody. Cytoplasm, cytoskeleton, microtubule organizing center, spindle pole body Note=Localizes to the centrosomes throughout interphase but migrates to the mitotic apparatus, including spindle pole bodies, mitotic spindle, and midbody, during mitosis.


Tissue Location

Expressed in all adult tissues examined except for lung and kidney.

Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human LATS1. The exact sequence is proprietary.

Images



Western blot analysis of LATS1 expression in LO2 (A), SGC7901 (B), HEK293T (C), A549 (D), mouse embryo (E), rat heart (F) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.